

(19)  **Canadian
Intellectual Property
Office**

An Agency of
Industry Canada

**Office de la Propriété,
Intellectuelle
du Canada**

Un organisme
d'Industrie Canada

(11) **CA 2 478 205** (13) **A1**

(40) 12.09.2003

(43) 12.09.2003

(12)

(21) 2 478 205

(22) 06.03.2003

(51) Int. Cl. 7: **C12N 15/09, C12P 21/02**

(85) 03.09.2004

(86) PCT/JP03/002675

(87) WO03/074699

(30) 2002-60374 JP 06.03.2002
2002-268726 JP 13.09.2002

(71) **TORAY INDUSTRIES, INC.,**
2-1 Nihonbashi-Muromachi 2-chome
Chuo-ku
103-8666, TOKYO, XX (JP).
**NATIONAL INSTITUTE OF AGROBIOLOGICAL
SCIENCES,**
1-2, Kannondai 2-chome,

(72) **Tsukuba-shi**
305-8602, IBARAKI, XX (JP).

(72) **TAMURA, TOSHIKI (JP).**
HIRAMATSU, SHINGO (JP).
TANAKA, TAKASHI (JP).
YAMADA, KATSUSHIGE (JP).

(74) **SMART & BIGGAR**

(54) **PROCEDE DE PRODUCTION DE PROTEINE PHYSIOLOGIQUEMENT ACTIVE UTILISANT UN VER A SOIE
GENETIQUEMENT MODIFIE**

(54) **PROCESS FOR PRODUCING PHYSIOLOGICALLY ACTIVE PROTEIN USING GENETICALLY MODIFIED
SILKWORM**

(57)

It is intended to provide a genetic engineering material for insects whereby a target protein can be easily purified without resort to a recombinant baculovirus and a process for producing a foreign protein with the use of this genetic engineering material. A genetically modified silk worm is obtained by transferring a foreign protein gene (for example, a cytokine gene) ligated to a promoter functioning in the silk gland into silkworm chromosome. Then the foreign protein (for example, cytokine) is extracted and purified from the silk gland or cocoon of the silkworm or its offspring. By transferring a gene cassette for expression wherein the 5'-terminal DNA sequence of fibroin H-chain gene is fused with the 3'-terminal DNA sequence of a foreign protein gene into silk gland cells or the like, the foreign protein can be produced in a large amount both inside and outside the silk gland cells as well as in silk yarn and cocoon.



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2478205 A1 2003/09/12

(21) 2 478 205

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) A1

(86) Date de dépôt PCT/PCT Filing Date: 2003/03/06

(87) Date publication PCT/PCT Publication Date: 2003/09/12

(85) Entrée phase nationale/National Entry: 2004/09/03

(86) N° demande PCT/PCT Application No.: JP 2003/002675

(87) N° publication PCT/PCT Publication No.: 2003/074699

(30) Priorités/Priorities: 2002/03/06 (2002-60374) JP;
2002/09/13 (2002-268726) JP

(51) Cl.Int.⁷/Int.Cl.⁷ C12N 15/09, C12P 21/02

(71) Demandeurs/Applicants:
TORAY INDUSTRIES, INC.; JP;
NATIONAL INSTITUTE OF AGROBIOLOGICAL
SCIENCES, JP

(72) Inventeurs/Inventors:
HIRAMATSU, SHINGO, JP;
TANAKA, TAKASHI, JP;
YAMADA, KATSUSHIGE, JP;
TAMURA, TOSHIKI, JP

(74) Agent: SMART & BIGGAR

(54) Titre : PROCÉDE DE PRODUCTION DE PROTEINE PHYSIOLOGIQUEMENT ACTIVE UTILISANT UN VER A SOIE
GENETIQUEMENT MODIFIE

(54) Title: PROCESS FOR PRODUCING PHYSIOLOGICALLY ACTIVE PROTEIN USING GENETICALLY MODIFIED
SILKWORM

(57) Abrégé/Abstract:

It is intended to provide a genetic engineering material for insects whereby a target protein can be easily purified without resort to a recombinant baculovirus and a process for producing a foreign protein with the use of this genetic engineering material. A genetically modified silk worm is obtained by transferring a foreign protein gene (for example, a cytokine gene) ligated to a promoter functioning in the silk gland into silkworm chromosome. Then the foreign protein (for example, cytokine) is extracted and purified from the silk gland or cocoon of the silkworm or its offspring. By transferring a gene cassette for expression wherein the 5' -terminal DNA sequence of fibroin H-chain gene is fused with the 3' -terminal DNA sequence of a foreign protein gene into silk gland cells or the like, the foreign protein can be produced in a large amount both inside and outside the silk gland cells as well as in silk yarn and cocoon.

Canada

<http://opic.gc.ca> • Ottawa-Hull K1A 0C9 • <http://cipo.gc.ca>

OPIC • CIPQ 191

OPIC



CIPQ

- 60 -

ABSTRACT

The present invention provides a genetic engineering material for insects that enables a target protein to be
5 purified easily, without requiring the use of recombinant baculovirus, while simultaneously providing a process for producing exogenous protein using that genetic engineering material. A gene recombinant silkworm is obtained by inserting an exogenous protein gene such as a
10 cytokine gene coupled to a promoter that functions in silk glands into a silkworm chromosome. An exogenous protein such as a cytokine is then extracted and purified from the silk glands or cocoon of that silkworm or its offspring. A large amount of exogenous protein can be
15 produced within silk gland cells, outside silk gland cells or in silk thread or a cocoon by inserting an expression gene cassette, in which the DNA sequence of the 3' terminal portion and the DNA sequence of the 5' terminal portion of fibroin H chain gene are fused to the
20 exogenous protein gene, into silk gland cells and so forth.

DESCRIPTION

PRODUCTION OF PHYSIOLOGICALLY ACTIVE PROTEINS USING GENE
RECOMBINANT SILKWORMS

5

Technical Field

The present invention relates to a process for producing a recombinant cytokine using a silkworm incorporating a cytokine gene in its chromosomes. In addition, the present invention relates to a gene recombinant silkworm having the property of producing a recombinant cytokine in a silk gland or cocoon and silk thread, and a vector for inserting an exogenous gene into silkworm chromosomes for producing the recombinant silkworm. In addition, the present invention also relates to a process for producing exogenous protein using insect cells, insect tissue or insects to which a gene has been inserted using the aforementioned vector. Moreover, the present invention relates to silk thread containing an exogenous protein produced by a recombinant silkworm obtained in the present invention.

Background Art

The production of exogenous proteins using gene recombination technology is used in various industries. The hosts used for their production consist mainly of *E. coli*, yeast, animal cells, plant cells and insect cells. However, a host has yet to be developed that is capable of efficiently producing all kinds of exogenous proteins and, as it is necessary to construct a production system for each target protein, a technical breakthrough is being sought in technology for producing exogenous proteins in individual hosts.

Systems using bacteria like *E. coli* or yeast have problems with posttranslational modification, and in some cases these systems are unable to synthesize proteins in a form that allows them to function adequately. In

- 2 -

addition, although animal cells allow proteins to be synthesized in a functional form, it is typically difficult to grow these cells and the production volume is low thereby making this uneconomical.

5 On the other hand, in the production of gene recombinant proteins using insects or insect cells, useful proteins having enzymatic or physiological activity can be produced comparatively inexpensively and modifications can be obtained, following protein
10 translation, that resemble those in animals. More specifically, a method in which a baculovirus incorporated a recombinant exogenous gene is infected into insects or insect cells allows the exogenous protein to be produced comparatively inexpensively, and
15 physiologically active proteins are known that have been commercialized as pharmaceuticals (Japanese Unexamined Patent Publication Nos. 61-9288 and 61-9297).

 In the case of the production of cytokines, which are physiologically active substances having
20 immunoregulatory functions and which are attracting attention in pharmaceutical applications, methods are disclosed in Japanese Unexamined Patent Publication Nos. 3-139276 and 9-234085 in which silkworms are inoculated with BmNPV containing a feline interferon- ω gene and a
25 canine interferon- γ gene, respectively. In addition, a process for producing human collagen using insect cells infected with baculovirus is known as an example of the production of a protein other than interferon using
insects (Japanese Unexamined Patent Publication No. 8-
30 23979).

 However, as technologies for producing recombinant proteins using insects or insect cells of the prior art use a recombinant virus to incorporate an exogenous gene, there is the problem of the need for deactivation or
35 containment of recombinant virus from the viewpoint of safety. In addition, in methods in which a recombinant

- 3 -

virus is inoculated into a silkworm, as the inoculation of the recombinant virus is troublesome task and the target exogenous protein is produced in silkworm hemolymph, it is necessary to purify the target recombinant protein from the large amount of contaminating proteins originating in the body fluids of the silkworm. Consequently, there was the problem of it being difficult to obtain a highly pure recombinant protein.

On the other hand, attempts have been made in recent years to recombine exogenous genes into insect chromosomes, and a method has been developed that uses homologous recombination to introduce and express in silkworm chromosomes a fused gene in which jellyfish green fluorescence protein gene was coupled to silkworm fibroin L chain gene using DNA of Autographa californica nuclear polyhedrosis virus (AcNPV), which is a type of nuclear polyhedrosis virus (Genes Dev., 13, 511-516, 1999), and a silkworm containing human collagen gene and a production process have been disclosed that utilize this technology (Japanese Unexamined Patent Publication No. 2001- 161214). Recently, research has been conducted on a method for expressing a protein encoded by an exogenous gene by stably introducing that exogenous gene into silkworm chromosomes using piggyBac, which is a transposon originating in a lepidopteron, using the jellyfish green fluorescence protein as a model, and the gene has been confirmed to be stably propagated to offspring by mating (Nature Biotechnology, 18, 81-84, 2000).

However, in the aforementioned method for inserting an exogenous protein gene into insect chromosomes using AcNPV, as a recombinant baculovirus (AcNPV) is used, there is still the problem of having to deactivate and contain the recombinant virus. In the example that used the piggyBac transposon, as the amount of green fluorescence protein produced is inadequate and as it is

- 4 -

also produced throughout the silkworm, sophisticated purification technology is required to recover the expressed recombinant green fluorescence protein in a highly pure form, thereby resulting in the method being uneconomical. In addition, the amount of recombinant protein produced is inadequate and extremely low.

Namely, in this technology for producing an exogenous protein using insect cells as a host, there are several problems such as the need to deactivate and contain the recombinant baculovirus, the difficulty in purifying the target protein from body fluid in which a large amount of contaminating proteins are present, as in the case of using silkworms, and the expressed amount of the target protein being low.

There are no known examples, thus far, of expressing a target protein by inserting a gene that encodes a physiologically active protein such as a cytokine gene into silkworm chromosomes. In addition, there are also no examples of having recovered a recombinant cytokine from a site, other than silkworm body fluid, such as a silk gland or silk thread secreted by silkworms, and confirming the physiological activity of the resulting cytokine. In addition, there are also no precedents regarding a silkworm capable of inheriting such properties. In addition, there are no examples of having produced a large amount of recombinant protein in silk thread using a recombinant silkworm produced using a transposon.

DISCLOSURE OF THE INVENTION

Although extensive research has been conducted on technologies for producing recombinant proteins using insects, there are problems such as the need to deactivate and contain the recombinant baculovirus in which the exogenous protein gene has been incorporated, or the need to take a lot of time and labor associated with inoculating the recombinant virus. In addition, the

production of exogenous protein in silkworms using recombinant baculovirus had the problem of it being difficult to extract and purify the target protein from body fluid containing large amounts of contaminating proteins.

Although studies have been conducted on technologies, for producing recombinant proteins, in which an exogenous protein gene has been inserted into silkworm chromosomes, these have problems consisting of the small amount of target exogenous protein produced and the difficulty in purifying the target protein from silkworm body fluid.

In consideration of these circumstances, the object of the present invention is to provide a genetic engineering material for insects that does not require the use of recombinant baculovirus and enables a target protein having physiological activity to be purified easily, while simultaneously providing a process for producing exogenous protein using that genetic engineering material.

As a result of extensive studies, the inventors of the present invention found that, by inserting a DNA sequence having a structure in which a gene that encodes a target protein is coupled downstream from a promoter specifically expressed in silkworm silk glands into silkworm chromosomes using DNA originating in a transposon, the target protein is produced in the silk glands, or the cocoon and the silk thread, in a form that retains physiological activity, thereby leading to completion of the present invention. In the present invention, as the recombinant protein can be recovered from the silk glands or silk and cocoon thread without containing a large amount of contaminants, it offers the advantage of allowing the target protein to be purified easily. Moreover, as a virus like baculovirus is not used, virus deactivation is not necessary thereby allowing the recombinant protein to be produced both

easily and safely.

In addition, as a result of conducting extensive studies focusing on the fact that the silkworm silk glands, and particularly the posterior silk gland, produces a large amount of fibroin that accounts for 70 to 80% of silk protein, and that the fibroin is secreted by the silk gland cells, the inventors of the present invention found that the amount of exogenous protein produced is increased considerably by inserting into silk gland cells a gene cassette in which the 5' terminal of an exogenous protein gene is coupled to the 3' terminal of a fibroin H chain gene 5' terminal portion containing a first intron of fibroin H chain gene downstream from a promoter expressed in the silk glands so that the amino acid frames are continuous. In addition, it was also found that a large amount of exogenous protein is secreted and produced by silk gland cells when a fused gene in which the 3' terminal portion of fibroin H chain gene is coupled to the 3' side of an exogenous protein gene, so that the amino acid frames are continuous, is expressed under the control of a promoter expressed in silk glands. In addition, it was also found that a recombinant silkworm produces a large amount of a target protein in its silk threads when a gene cassette was produced in which a DNA sequence of the 5' terminal portion containing a first intron of fibroin H chain gene on the 5' side of an exogenous protein gene, and a DNA sequence of the 3' terminal portion of fibroin H chain gene on the 3' side, were respectively designed so that their amino acid frames were continuous, followed by producing a recombinant silkworm in which that gene cassette was inserted into its chromosomes.

The inventors of the present invention succeeded in producing a large amount of exogenous protein in silk gland cells, outside silk gland cells and in silk thread by inserting into silk gland cells and so forth an expression gene cassette in which the DNA sequence of the

- 7 -

5' terminal portion and the DNA sequence of the 3' portion of fibroin H chain gene were fused to an exogenous protein gene, and were able to establish an exogenous protein production technology that facilitates purification by producing an exogenous protein using silk glands instead of using a recombinant baculovirus.

Namely, the present invention relates to a process for producing a recombinant cytokine comprising producing a gene recombinant silkworm that incorporates a cytokine gene in its chromosomes, and recovering the cytokine from the silk glands or cocoon and silk thread. Moreover, the present invention also relates to a gene recombinant silkworm in which a cytokine gene is incorporated, and a gene recombinant vector used to insert the cytokine gene into the silkworm.

Moreover, the present invention relates to a genetic engineering material, such as the gene cassette or vector described below, capable of being used for exogenous protein production in insects, a transformant, a process for producing exogenous protein using that transformant, and silk thread containing exogenous protein.

Thus, the present invention provides 1) a gene cassette for expressing an exogenous protein comprising (1) a promoter expressed in silk glands, and (2) a gene coupled downstream from (1) in which the 5' terminal portion of fibroin H chain gene is fused to the 5' side of an exogenous protein structural gene.

Moreover, the present invention provides 2) a gene cassette for expressing an exogenous protein comprising (1) a promoter expressed in silk glands, and (2) a gene coupled downstream from (1) in which the 3' terminal portion of fibroin H chain gene is fused to the 3' side of an exogenous protein structural gene not containing a stop codon. Alternatively, the present invention provides a gene cassette for expressing an exogenous protein comprising (1) a promoter expressed in silk glands, and (2) a gene coupled downstream from (1) in

- 8 -

which an exogenous protein structural gene is fused to the 3' side of the 3' terminal portion of fibroin H chain gene.

Moreover, the present invention provides 3) a gene cassette for expressing an exogenous protein comprising (1) a promoter expressed in silk glands, and (2) a gene coupled downstream from (1) in which the 5' terminal portion of fibroin H chain gene is fused to the 5' side of an exogenous protein structural gene not containing a stop codon, and the 3' terminal portion of fibroin H chain gene is fused to the 3' side of the structural gene.

In addition, the present invention provides 4) an expression vector for insect cells containing a gene cassette for expressing an exogenous protein according to any of the aforementioned 1) through 3).

Moreover, the present invention provides 5) a process for producing exogenous protein comprising inserting an expression vector for insect cells according to the aforementioned 4) into insect cells.

Moreover, the present invention provides 6) a process for producing exogenous protein comprising producing a recombinant silkworm in which a gene cassette for expressing an exogenous protein according to any of the aforementioned 1) through 3) is incorporated in its chromosomes, and after producing the exogenous protein in the silk glands or silk thread of the resulting recombinant silkworm, recovering the exogenous protein from the silk glands or silk thread.

In addition, the present invention provides 7) a recombinant silkworm in which a gene cassette for expressing an exogenous protein according to any of the aforementioned 1) through 3) is incorporated in its chromosomes, and has the ability to produce exogenous protein in its silk glands or silk thread.

Moreover, the present invention provides 8) a silk thread containing an exogenous protein produced by the

silkworm according to the aforementioned 7).

BRIEF DESCRIPTION OF THE DRAWINGS

5 Fig. 1 is a drawing showing a restriction map of gene insertion vector pigSIB.

Fig. 2 is a drawing showing a restriction map of gene insertion vector pigFIB.

Fig. 3 is a drawing showing a restriction map of plasmid pHA3PIG having a transposase.

10 Fig. 4 is a drawing showing the results of treating the genomic DNA of 11 silkworms (G1) obtained from the positive moth groups of Table 1 with EcoRV and XmnI, followed by performing Southern blotting analysis using a feline interferon- ω gene as a probe.

15 Fig. 5 is a drawing showing the antiviral activity of a silk thread extract of a recombinant silkworm into which was inserted feline interferon- ω gene coupled to fibroin H chain promoter. The sample in the dyed lane is shown to have activity.

20 Fig. 6 is a drawing showing the results of treating silkworm silk gland genomic DNA obtained from the positive moth groups of Table 3 (3 moth groups from Experiment 1 and 2 moth groups from Experiment 2) with EcoRI or BglIII followed by performing Southern blotting analysis using feline interferon- ω gene as a probe.

25 Fig. 7 is a drawing of the detection of expression of feline interferon mRNA in a middle silk thread of a gene recombinant silkworm by RT-PCR.

30 Fig. 8 is a drawing showing the antiviral activity of a middle silk gland extract and a cocoon and silk thread extract of a recombinant silkworm into which was inserted feline interferon- ω gene coupled to a sericin promoter. The sample in the dyed lane is shown to have activity.

35 Fig. 9 is a drawing showing the procedure for producing a construct for gene insertion containing a P-

- 10 -

IC-A gene cassette (first half).

Fig. 10 is a drawing showing the procedure for producing a construct for gene insertion containing a P-IC-A gene cassette (second half).

5 Fig. 11 is a drawing showing the procedure for producing a construct for gene insertion containing an HP-IC-HA gene cassette (first half).

10 Fig. 12 is a drawing showing the procedure for producing a construct for gene insertion containing an HP-IC-HA gene cassette (second half).

Fig. 13 is a drawing showing the procedure for producing a construct for gene insertion containing an HUP-IC-HA gene cassette (first half).

15 Fig. 14 is a drawing showing the procedure for producing a construct for gene insertion containing an HUP-IC-HA gene cassette (second half).

Fig. 15 is a drawing showing the procedure for producing a construct for gene insertion containing an HP-IC-A gene cassette (first half).

20 Fig. 16 is a drawing showing the procedure for producing a construct for gene insertion containing an HP-IC-A gene cassette (second half).

25 Fig. 17 is a drawing of the analysis of the expression of β -galactosidase in cultured silkworm silk glands by Western analysis. The first exon, first intron and second exon regions of fibroin H chain gene were clearly determined to play an important role in synthesis or gene expression of proteins within cells. In addition, secretion outside the cells was also confirmed.

30 Fig. 18 is a drawing of the analysis of the expression of recombinant protein in silkworm silk gland tissue by Western analysis. The first exon, first intron and second exon regions of fibroin H chain gene were reconfirmed to play an important role in dramatically
35 improving the expression of recombinant protein in silkworm posterior silk gland cells. In addition, a gene region that improves the amount of protein produced was

- 11 -

found in a roughly 5.5 kbp upstream region from the fibroin promoter.

Fig. 19 is a drawing of the analysis of the production of recombinant protein in silk thread by Western analysis. The 3' terminal portion of fibroin H chain gene was determined to play an important role in secretion of protein synthesized within silk gland cells to silk thread. In addition, a gene region that improves the amount of protein produced was reconfirmed in the roughly 5.5 kbp upstream region from the fibroin promoter.

BEST MODE FOR CARRYING OUT THE INVENTION

Cytokines are proteins produced by various cells that have immunoregulatory activity, antiviral activity and blood cell growth activity on hematopoietic cells and immunocytes. Their activity is demonstrated as a result of forming a precise higher order structure and bonding to specific receptors on the cell membrane. They have previously been applied clinically to humans and animals based on the characteristics of their activity.

Although there are no particular limitations on the cytokines of the present invention, they should be cytokines for which their physiological activity is maintained when expressed in silkworms, examples of which include physiologically active substances having immunoregulatory activity, antiviral activity or blood cell growth activity and so forth such as human interferon- α , β and γ (J. Interferon Res. 5, 521-526, 1985; Nucleic Acids Res. 10, 2487-2501, 1982), human interleukin-12 (J. Immunol. 146, 3074-3081, 1991), human granulocyte colony stimulating factor (Nature, 319, 415-418, 1986), human erythropoietin (Nature, 313, 806-810, 1985), human thrombopoietin (Cell, 77, 1177-1124, 1994), feline interferon- ω , (Biosci. Biotech. Biochem., 56, 211-214, 1992, GenBank database registration no. E04599),

- 12 -

feline erythropoietin (GenBank database registration no. FDU00685), feline granulocyte colony stimulating factor (Gene, 274, 263-269, 2001), canine interferon- γ (GenBank database registration no. S41201), canine interleukin-12 (Japanese Unexamined Patent Publication No. 10-36397) and
5 canine granulocyte colony stimulating factor (US Patent No. 5606024). Preferable examples of cytokines include interferons and colony stimulating factors, and these preferably include interferon- α , β , γ , ω and τ along with
10 colony stimulating factor, erythropoietin and thrombopoietin. More preferable examples of cytokines include feline interferon- ω , feline granulocyte colony stimulating factor and human interferon- β .

Feline interferon- ω gene is obtained by cutting out
15 from a plasmid extracted from, for example, *E. coli* (pFeIFN1) (Patent Microorganism Depository No. 1633). In addition, this gene can also be obtained from rBNV100 produced by co-transfecting into established silkworm cells with a recombinant plasmid produced by ligating
20 feline interferon- ω gene to a silkworm cloning vector (T. Horiuchi, et al., Agric. Biol. Chem., 51, 1573-1580, 1987), and silkworm nuclear polyhedrosis virus.

Feline granulocyte colony stimulating factor can be obtained by stimulating CRFK cells, which are cultured
25 cells originating in feline kidney, with LPS followed by recovering mRNA from the cells and then carrying out PCR using the cDNA obtained by reverse transcription as a template and using primers established with reference to GenBank database registration no. AB042552.

30 Human interferon- β gene can be acquired by cutting out from plasmid ORF-hIFN- β (Invitrogen) that encodes its cDNA.

The method for inserting a gene into silkworm chromosomes used in the present invention should enable
35 the gene to be stably incorporated and expressed in the

chromosomes, and be stably propagated to offspring, as well, by mating. Although a method using micro-injection into silkworm eggs or a method using a gene gun can be used, a method that is used preferably consists of the
5 micro-injection into silkworm eggs with a target gene containing vector for insertion of an exogenous gene into silkworm chromosomes and helper plasmid containing a transposon gene (Nature Biotechnology 18, 81-84, 2000) simultaneously.

10 The target gene is inserted into reproductive cells in a recombinant silkworm that has been hatched and grown from the micro-injected silkworm eggs. Offspring of a recombinant silkworm obtained in this manner are able to stably retain the target gene in their chromosomes. The
15 gene recombinant silkworm obtained in the present invention can be maintained in the same manner as ordinary silkworms. Namely, up to fifth instar silkworms can be raised by incubating the eggs under normal conditions, collecting the hatched larva to artificial
20 feed and then raising under the same conditions as ordinary silkworms.

Gene recombinant silkworms obtained in the present invention are able to pupate and produce a cocoon in the same manner as ordinary silkworms. Males and females are
25 distinguished in the pupa stage, and after having transformed into moths, males and females mate and eggs are gathered on the following day. The eggs can be stored in the same manner as ordinary silkworm eggs. The gene recombinant silkworms of the present invention can
30 be maintained on subsequent generations by repeating the breeding as described above, and can be increased to large numbers.

The exogenous gene insertion vector used for the purpose of inserting a cytokine gene used in the present
35 invention into silkworm chromosomes is not subject to any particular limitations provided it is designed so as to precisely control cytokine expression. Normally, it has

- 14 -

a structure in which the cytokine gene is coupled to downstream from a promoter specifically expressed in the silk glands and upstream from an arbitrary poly A sequence, and has a pair of DNA sequences originating in a transposon outside these gene sequences. Moreover, a signal sequence originating in an arbitrary gene may be coupled between the cytokine gene and the promoter, and an arbitrary gene sequence may also be coupled between the cytokine gene and poly A. In addition, an artificially designed and synthesized gene sequence can also be coupled. In addition, a sequence for replication within a bacterial host, antibiotic resistance gene, fluorescence protein gene or LacZ gene and so forth can also be coupled as necessary. For example, the gene of green fluorescence protein GFP coupled downstream from a suitable promoter can be inserted at a suitable location between a pair of transposon DNA sequences. As a result, this facilitates screening for gene recombinant silkworms. In addition, this vector may also contain all or a portion of pUC9, pUC19 or other plasmids originating in *E. coli*.

Moreover, although there are no particular limitations on the promoter used here, and any promoter originating in any organism can be used provided it acts effectively within silkworm cells, a promoter that has been designed to specifically induce protein in silkworm silk glands is preferable. Examples of silkworm silk gland protein promoters include fibroin H chain promoter, fibroin L chain promoter, p25 promoter and sericin promoter.

Examples of other gene sequences used in addition to the promoter include signal sequences, poly A sequences and other sequences that control gene expression. These are not limited to specific gene sequences, but rather those which are suitable for expression of the target gene can be selected. Examples include sequences originating in the target protein such as signal

- 15 -

sequences of cytokines such as feline interferon- ω and poly A sequences, and signal sequences and poly A sequences of insect protein contained in the silkworm serving as the host. Alternatively, other examples
5 include sequences that have been proven to be generally effective for expressing proteins such as SV40 poly A and bovine growth hormone poly A. By changing the gene sequence of the aforementioned promoters and the other sequences coupled with the cytokine genes, the locations
10 where they are expressed and the amounts expressed can be controlled.

In the present invention, a "gene cassette for expressing an exogenous protein" refers to a set of DNA required for a synthesis of protein encoded by the
15 exogenous protein structural gene in the case of being inserted into insect cells. This gene cassette for expressing an exogenous protein contains an exogenous protein structural gene and a promoter that promotes expression of that gene. Normally, it also contains a
20 terminator and poly A addition region, and preferably contains a promoter, exogenous protein structural gene, terminator and poly A addition region. Moreover, it may also contain a secretion signal gene coupled between the promoter and the exogenous protein structural gene. An
25 arbitrary gene sequence may also be coupled between the poly A addition sequence and the exogenous protein structural gene. In addition, an artificially designed and synthesized gene sequence can also be coupled.

In addition, a "gene cassette for inserting a gene"
30 refers to a gene cassette for expressing an exogenous gene having an inverted repetitive sequence of a pair of piggyBac transposons on both sides, and consisting of a set of DNA inserted into insect cell chromosomes through the action of the piggyBac transposons.

35 There are no particular limitations on the method used to acquire DNA used in the present invention. Examples of such methods include a method in which a

- 16 -

required gene region is amplified and acquired using a polymerase chain reaction (PCR) based on known genetic information, and a method in which a genome library or cDNA library is screened using homology as an indicator based on known genetic information. In the present invention, these genes include variants resulting from genetic polymorphism and artificial mutation treatment using mutagens and so forth. Genetic polymorphism refers to that in which a portion of the base sequence of a gene is altered by a sudden spontaneous mutation in the gene.

Although there are no particular limitations on the promoter in the gene cassette for expressing an exogenous protein, that having a high level of activity that promotes expression of an exogenous protein gene is preferable. Although examples include the promoter of drosophila heat shock protein gene described in Japanese Unexamined Patent Publication No. 6-261770 and the promoter of silkworm actin gene (Nature Biotechnology 18, 81-84, 2000), promoters having a high level of promoting activity in silkworm silk gland cells are preferable, examples of which include the promoters of fibroin H chain gene (base numbers 255-574 of GenBank registration no. V00094), fibroin L chain gene (Gene, 100, 151-158; GenBank registration no. M76430) and sericin gene (base numbers 599-1656 of GenBank registration no. AB007831).

"Exogenous protein structural gene" refers to a gene not possessed by host cells in which a gene is to be expressed, and which encodes a protein not inherently produced by the host cells. Although there are no particular limitations thereon, in consideration of industrial value, examples include genes of proteins that are produced by humans or mammals such as genes of growth hormones, cytokines, growth factors and cell structural proteins. In addition, genes of enzymes and various proteins produced by microbes, plants or insects are also included in the scope of the present invention.

In the gene cassette for expressing an exogenous

- 17 -

protein in the present invention, the 5' terminal portion of fibroin H chain gene is a DNA sequence having action that enhances expression of exogenous protein gene by a promoter, and contains a first exon of fibroin H chain gene, all or a portion of a first intron, and a portion of a second exon. By fusing the 5' side of an exogenous protein structural gene to the 3' side of this second exon so that the amino acid reading frame is contiguous, the amount of exogenous protein produced can be improved. However, since surplus amino acid residues are added to the N terminal side of the target exogenous protein if the second exon portion is too long, there are cases in which the structure or activity of the target exogenous protein is lost. Consequently, it is necessary that the second exon portion have a suitable length according to the purpose. In many cases, favorable results can be obtained by making the second exon portion to extend to immediately after or up to several amino acid residues from the secretion signal gene of fibroin H chain gene. In addition, as the region upstream from the 5' side of fibroin H chain gene promoter, namely a roughly 5.5 kbp upstream region, is considered to be the region that enhances promoter activity, adding this region can be expected to increase the amount of target protein expressed.

In the case of producing an exogenous protein in silkworm silk glands, the 3' terminal portion of fibroin H chain gene is a DNA sequence having the effect of causing secretion of a large amount of exogenous protein outside the silk gland cells. A recombinant silkworm in which a gene cassette for expressing an exogenous protein, in which the 3' terminal portion of fibroin H chain gene serving as the signal for secreting into silk thread is fused to the 3' side, is inserted into its chromosomes is able to produce exogenous protein in its silk thread. In addition, the 3' terminal portion of fibroin H chain gene may be present upstream or

downstream from the exogenous protein gene or within the exogenous protein gene.

5 In the case at least one cysteine residue is present in this portion and the 3' terminal of fibroin H chain gene is used as is, the cysteine residue is located at the 20th residue from the carboxyl terminal of the fibroin H chain gene. This cysteine fulfills the role of bonding to fibroin L chain by a disulfide bond. There are no particular limitations on the length of the DNA sequence of the 3' terminal portion of fibroin H chain gene provided it does not inhibit formation of the disulfide bond with fibroin L chain. As a repetitive DNA sequence continues from about 100 or more bases upstream from the 3' terminal of fibroin H chain, cleaving the DNA sequence of this upstream portion to an arbitrary length is difficult with a restriction endonuclease. Thus, in consideration of the ease of genetic engineering techniques, roughly 100 base pairs on the 3' portion where the repetitive DNA sequence of fibroin H chain gene ends can be preferably used for the 3' terminal portion of fibroin H chain. In addition, as a large number of amino acids originating in the carboxyl terminal of fibroin H chain protein bond to the carboxyl terminal or amino terminal of the exogenous protein if the 3' terminal portion of fibroin H chain gene is excessively long, there are cases in which the structure or activity of the target exogenous protein is lost. Thus, there are cases in which it is necessary to make the DNA sequence of the 3' terminal portion of fibroin H chain gene as short as possible depending on the target protein.

30 Although there are no particular limitations on the poly A region, a poly A region of a protein gene expressed in large amounts in silk glands, such as fibroin H chain, fibroin L chain or sericin, can be used preferably.

35 A vector in the present invention refers to that having a cyclic or linear DNA structure. A vector

capable of replicating in *E. coli* and having a cyclic DNA structure is particularly preferable. This vector can also incorporate a marker gene such as an antibiotic resistance gene or jellyfish green fluorescence protein gene for the purpose of facilitating selection of transformants.

Although there are no particular limitations on the insect cells used in the present invention, they are preferably lepidopteron cells, more preferably *Bombyx mori* cells, and even more preferably silkworm silk gland cells or cells contained in *Bombyx mori* eggs. In the case of silk gland cells, posterior silk gland cells of fifth instar silkworm larva are preferable because there is active synthesis of fibroin protein and they are easily handled.

There are no particular limitations on the method used to incorporate a gene cassette for expression of exogenous protein and a vector into the insect cells. Although the calcium phosphate method, methods using electroporation, methods using liposomes, methods using a gene gun and methods using micro-injection can be used for incorporation into cultured insect cells, in the case of incorporating into silkworm silk gland cells, for example, a gene can be easily incorporated into posterior silk gland tissue removed from the body of a fifth instar silkworm larvae using a gene gun.

Gene incorporation into the posterior silk gland using a gene gun can be carried out by, for example, bombarding gold particles coated with a vector containing a gene cassette for expressing exogenous protein into a posterior silk gland immobilized on an agar plate and so forth using a particle gun (Bio-Rad, Model No. PDS-1000/He) at an He gas pressure of 1,100 to 1,800 psi.

In the case of incorporating a gene into cells contained in eggs of *Bombyx mori*, a method using micro-injection is preferable. Here, in the case of performing micro-injection into eggs, it is not necessary to micro-

- 20 -

inject into the cells of the eggs directly, but rather a gene can be incorporated by simply micro-injecting into the eggs.

5 A recombinant silkworm containing the "gene cassette for expressing an exogenous protein" of the present invention in its chromosomes can be acquired by micro-injecting a vector having a "gene cassette for inserting a gene" into the eggs of *Bombyx mori*. For example, a first generation (G1) silkworm is obtained by
10 simultaneously micro-injecting a vector having a "gene cassette for inserting a gene" and a plasmid in which a piggyBac transposase gene is arranged under the control of silkworm actin promoter into *Bombyx mori* eggs according to the method of Tamara, et al. (Nature
15 Biotechnology 18, 81-84, 2000), followed by breeding the hatched larva and crossing the resulting adult insects (G0) within the same group. Recombinant silkworms normally appear at a frequency of 1 to 2% among this G1 generation.

20 Selection of recombinant silkworms can be carried by PCR using primers designed based on the exogenous protein gene sequence after isolating DNA from the G1 generation silkworm tissue. Alternatively, recombinant silkworms can be easily selected by inserting a gene encoding green
25 fluorescence protein coupled downstream from a promoter capable of being expressed in silkworm cells into a "gene cassette for inserting a gene" in advance, and then selecting those individuals that emit green fluorescence under ultraviolet light among G1 generation silkworms at
30 first instar stage.

In addition, in the case of the micro-injection of a vector having a "gene cassette for inserting a gene" into *Bombyx mori* eggs for the purpose of acquiring recombinant silkworms containing a "gene cassette for expressing an
35 exogenous protein" in their chromosomes, recombinant silkworms can be acquired in the same manner as described above by simultaneously micro-injecting a piggyBac

transposase protein.

A piggyBac transposon refers to a transfer factor of DNA having an inverted sequences of 13 base pairs on both ends and an ORF inside of about 2.1k base pairs.

5 Although there are no particular limitations on the piggyBac transposon used in the present invention, examples of those that can be used include those originating in *Trichoplusia ni* cell line TN-368, *Autographa californica* NPV (AcNPV) and *Galleria mellonea* NPV (GmMNPV). A piggyBac transposon having gene and DNA transfer activity can be preferably prepared using plasmids pHA3PIG and pPIGA3GFP having a portion of a piggyBac originating in *Trichoplusia ni* cell line TN-368 (Nature Biotechnology 18, 81-84, 2000).

15 The structure of the DNA sequence originating in a piggyBac is required to have a pair of inverted terminal sequences containing a TTAA sequence, and has an exogenous gene such as a cytokine gene inserted between those DNA sequences. It is more preferable to use a transposase in order to insert an exogenous gene into silkworm chromosomes using a DNA sequence originating in a transposon. For example, the frequency at which a gene is inserted into silkworm chromosomes can be improved considerably by simultaneously inserting DNA capable of expressing a piggyBac transposase to enable the transposase transcribed and translated in the silkworm cells to recognize the two pairs of inverted terminal sequences, cut out the gene fragment between them, and transfer it to silkworm chromosomes.

25 30 The gene recombinant silkworm used in the present invention refers to a silkworm which has had inserted into its chromosomes an exogenous protein gene, and after treating the silkworm chromosomal DNA with restriction endonuclease in accordance with ordinary methods, yields a positive signal when subjected to Southern blotting using the exogenous protein gene labeled in accordance with ordinary methods as a probe. There are no

- 22 -

particular limitations on the gene locus on the chromosome into which a cytokine gene has been inserted provided it is a site that does not inhibit silkworm development, differentiation and growth. The recombinant
5 silkworm has the ability to produce exogenous protein in its silk gland cells, silk gland lumen and silk thread. In addition, the recombinant silkworm is able to develop and mate normally, stably retain the inserted exogenous protein gene, and transmit that gene to its offspring.
10 Thus, the amount of exogenous protein produced can easily be increased by increasing the number of recombinant silkworms through crossing. Crossing between transgenic silkworm strain and non-transgenic strain can increase the amount of the produced exogenous protein. In this
15 case, it is necessary to cross the silkworms while suitably selecting those silkworms into which the target exogenous protein gene has been inserted. In this case, offspring that have inherited the gene of the recombinant silkworm can be easily evaluated by analyzing a marker
20 gene used to select the recombinant silkworms or the presence or structure of the exogenous protein gene by PCR or Southern blotting and so forth using cell DNA obtained from an arbitrary tissue.

Insect cells and silkworm silk glands containing the
25 gene cassette for expression of an exogenous protein of the present invention can produce exogenous protein in culture supernatant or their cells by respectively culturing in culture liquid suitable for their culturing. For example, BmN cells, which are silkworm ovary cells
30 that have been inserted with the expression gene cassette of the present invention, produce a target exogenous protein after 3 or 4 days of culturing by culturing at 27°C in TC-100 medium (PharMingen). In addition, silkworm posterior silk gland produces exogenous protein
35 by culturing at 25°C in Grace's insect medium after being excised aseptically from, for example, fifth instar larva. In the case of producing protein in silk glands,

- 23 -

it is preferable to maintain a high dissolved oxygen concentration in the medium, and culture while removing low molecular weight factors that inhibit protein synthesis that accumulate in the medium by, for example,
5 an ultrafiltration membrane since this allows protein synthesis to proceed for a long period of time.

A silk gland inserted with an exogenous protein gene fused to the 3' terminal of fibroin H chain gene of the present invention is capable of producing a large amount
10 of a target exogenous protein in culture supernatant. Since nearly all contaminating proteins in the silk gland culture supernatant are fibroin, the target protein can be easily purified from the silk gland culture supernatant, and as a result, a highly pure target
15 protein can be obtained.

The recombinant silkworm obtained in the present invention can be raised in the same manner as ordinary silkworms, and is able to produce exogenous protein by raising under ordinary conditions. The amount of
20 exogenous protein produced can be improved by optimizing the temperature, humidity and feeding conditions, etc. during the fifth instar period in particular corresponding to the target exogenous protein.

A recombinant silkworm inserted with an exogenous
25 protein gene fused to the 3' terminal of fibroin H chain gene of the present invention is able to produce a large amount of a target exogenous protein in its cocoon. The target exogenous protein can also be easily purified and recovered from the resulting cocoon. In addition,
30 depending on the function of the exogenous protein produced, silk thread containing the resulting exogenous protein can be used directly or in a partially processed form in various industrial applications.

Exogenous protein can be obtained from the silk
35 gland or cocoon and silk thread of a recombinant silkworm obtained in the present invention by a suitable extraction procedure. Although there are no particular

- 24 -

limitations on the solvent used to extract exogenous protein from silk glands or cocoon and silk thread, an aqueous solvent system is preferable in many cases. An aqueous solution used for extraction may contain a
5 suitable solute for promoting extraction of the exogenous protein, examples of which include inorganic acids such as phosphoric acid, organic acids such as acetic acid, citric acid and malic acid, salts such as sodium
10 chloride, urea, guanidine hydrochloride and calcium chloride, and polar organic solvents such as ethanol, methanol, acetonitrile and acetone. In addition, there are also no particular limitations on the pH of the extraction solution, and any arbitrary pH can be used provided it does not deactivate the function of the
15 target exogenous protein.

There are no particular limitations on the method for isolating and purifying the extracted exogenous protein, and ordinary protein purification methods can be used. For example, a target useful protein can be
20 purified and isolated by combining chromatography using a silica gel carrier, ion exchange carrier, gel permeation carrier, chelating carrier or pigment-loaded carrier and so forth, ultrafiltration, gel permeation, dialysis, desalting by salting out or concentration and so forth
25 using an inherently possessed function as an indicator. For example, feline interferon- ω can be recovered in the soluble fraction obtained by homogenizing silk glands or cocoon and silk thread of a silkworm into which has been inserted a feline interferon- ω gene with 20 mM phosphate
30 buffer (pH 7.0). Moreover, the purity of the feline interferon- ω can be increased by adsorbing the resulting extract liquid onto, for example, a Blue Sepharose carrier and eluting the resulting buffer solution containing the extract liquid after washing.
35 Cytokines produced in this manner can be used in pharmaceutical applications as well as various

- 25 -

measurement and diagnostic applications in the same manner as cytokines produced with other conventional production processes. In this case, they may also be used as a mixture to which various additives have been added. In addition, the tissue or cocoon and silk thread of a silkworm in which cytokines have been expressed can also be used directly or after processing as fibers for medical or clothing use. In addition, the tissue or silk thread of a recombinant silkworm in which enzymes have been expressed can be used directly in enzyme reactions.

Examples

Although the following provides a more detailed explanation of the present invention by indicating its examples, the present invention is not limited to the descriptions of these examples.

Reference Example - Method for Measuring Antiviral Activity

The physiological activity of interferon was measured according to the following method as antiviral activity.

Antiviral activity was measured by the CPE method using vesicular stomatitis virus (VSV) for the virus, and using feline Fc9 cells (J.K. Yamamoto, et al.: Vet. Immunol. and Immunopathol., 11, 1-19, 1986) for the susceptible cells in the case of feline interferon- ω , or human FL cells in the case of human interferon- β . Namely, a sample diluent was added to the uppermost row of 96-well microtiter plate in which susceptible cells cultured at 37°C to confluency, and then serially diluting in two-fold increments moving towards the lower end of the plate.

After culturing for 20 to 24 hours at 37°C, VSV was added followed by additionally culturing for 16 to 20 hours at 37°C. Viable susceptible cells adhered to the microtiter plate were then stained with crystal-violet

- 26 -

stain containing 20% formalin, and as a result of measuring the optical absorbance at 570 nm for the amount of crystal-violet remaining on the microplate, antiviral activity was determined by comparison with a standard.

5 Intercat (Toray) adjusted to 1000 units/ml with cell culturing medium was used for the standard for feline interferon- ω , while Feron (Toray) prepared to 1000 units/ml with cell culturing medium was used for the standard for human interferon- β . In addition, samples

10 were used for the measurement of antiviral activity after diluting 15-fold with cell culturing medium.

Example 1 - Preparation of *Bombyx mori* Genomic DNA

Fifth instar third day silkworms were dissected to remove posterior silk gland tissue. After washing with

15 1xSSC, 200 μ l of DNA extraction buffer (50 mM Tris-HCl (pH 8.0), 1 mM EDTA (pH 8.0), 100 mM NaCl) were added. After adding Proteinase K (final concentration: 20 μ g/ml) and adequately grinding up the tissue with a grinder, 350 μ l of DNA extraction buffer and 60 μ l of 10% SDS were

20 added followed by incubating for 2 hours at 50°C. After adding 500 μ l of Tris-HCl-saturated phenol (pH 8.0) and mixing for 10 minutes, the supernatant was recovered by centrifuging for 5 minutes at 4°C and 10,000 rpm. After adding an equal volume of phenol/chloroform/isoamyl

25 alcohol (25:24:1) to the supernatant and mixing, the resulting mixture was centrifuged. Phenol/chloroform/isoamyl alcohol was again added followed by centrifuging and recovery of the supernatant. After adding an equal volume of chloroform/isoamyl

30 alcohol (24:1) and mixing, the mixture was centrifuged. Chloroform/isoamyl alcohol was again added to the resulting supernatant followed by centrifuging and recovery of the supernatant. 1/10 volume 3 M sodium acetate (pH 5.2) was then added to the resulting

35 supernatant and mixed and, after additionally adding 2.5

- 27 -

volumes of cold ethanol and allowing it to stand undisturbed for 30 minutes at -80°C , the mixture was centrifuged for 10 minutes at 4°C and 15,000 rpm to precipitate genomic DNA. After washing the DNA precipitate with 70% ethanol, the precipitate was air-dried. The precipitate was then dissolved in sterile water containing RNase to 100 $\mu\text{g}/\text{ml}$ to prepare a diluted genomic DNA solution.

Example 2 - Gene Preparation

The genes used were acquired by PCR by producing primers for the sequences on both ends using known sequences and using suitable DNA sources for the templates. Restriction sites were added to the ends of the primers for the subsequent gene construction procedure.

Feline interferon- ω gene (base numbers 9-593 of GenBank registration no. S62636) was acquired by PCR using two types of primers consisting of primer 3 (SEQ. ID No. 3) and primer 4 (SEQ. ID No. 4) and using baculovirus rBNV100 encoding feline interferon- ω gene for the template. rBNV100 can be produced by, for example, cutting out FeIFN gene from a plasmid extracted from *E. coli* (pFeIFN1) (Patent Microorganism Depository No. 1633), coupling to a silkworm cloning vector (T. Horiuchi, et al., Agric. Biol. Chem., 51, 1573-1580, 1987), and co-transfecting silkworm established cells with the recombinant plasmid produced and silkworm nuclear polyhedrosis virus DNA.

Sericin-1 gene promoter (base numbers 599-1656 of GenBank registration no. AB007831) was acquired by PCR using two types of primers consisting of primer 5 (SEQ. ID No. 5) and primer 6 (SEQ. ID No. 6) and using silkworm chromosomal DNA for the template. Fibroin H chain gene promoter (base numbers 255-574 of GenBank registration no. V00094) was acquired by PCR using two types of primers consisting of primer 7 (SEQ. ID No. 7) and primer

- 28 -

8 (SEQ. ID No. 8) and using silkworm chromosomal DNA for the template. Bovine growth hormone gene poly A (pcDNA3.1(+)) sequence numbers 1011-1253) was acquired by PCR using two types of primers consisting of primer 9 (SEQ. ID No. 9) and primer 10 (SEQ. ID No. 10) and using plasmid pcDNA3.1(+) vector (Invitrogen) for the template.

PCR was carried out in accordance with the accompanying protocol using KODplus (Toyobo). Namely, after adding 10 ng of each template in the case of a plasmid or 100 ng in the case of chromosomal DNA, 30 pmol of each primer and 10 μ l of the 10xPCR buffer provided, each reagent was added to a concentration of 1 mM $MgCl_2$, 0.2 mM dNTPs and 2 units of KODplus followed by bringing up to a final volume of 100 μ l. The PCR components were then reacted for 30 cycles using a Perkin-Elmer DNA thermal cycler under DNA denaturation conditions of 94°C for 15 seconds, primer annealing conditions of 55°C for 30 seconds, and elongation conditions of 68°C for 30 to 60 seconds.

These reaction solutions were electrophoresed with 1 to 1.5% agarose gel, and DNA fragments consisting of a roughly 580 bp fragment in the case of feline interferon- ω gene, a roughly 1 kbp fragment in the case of sericin-1 promoter, a roughly 320 bp fragment in the case of fibroin H chain promoter, and a roughly 230 bp fragment in the case of bovine growth hormone poly A were extracted and prepared in accordance with ordinary methods. After phosphorylating these DNA fragments with polynucleotide kinase (Takara Shuzo), they were ligated to pUC19 vector subjected to dephosphorylation treatment after being cleaved with HincII by reacting overnight at 16°C using DNA Ligation Kit Ver. 2 (Takara Shuzo). These were then used to transform *E. coli* in accordance with ordinary methods and the resulting transformants were confirmed to contain the PCR fragments by performing PCR on the resulting colonies under the same conditions as

- 29 -

previously described to prepare plasmids in which the PCR fragments were inserted according to ordinary methods. These plasmids were sequenced to confirm that the resulting fragments consisted of the base sequences of each gene.

Example 3 - Production of Plasmids for Gene Insertion

pigA3GFP (Nature Biotechnology 18, 81-84, 2000) was used for the plasmid for gene insertion. Namely, vector pigA3GFP is a vector in which after removing a region encoding transposase from plasmid p3E1.2 disclosed in US Patent No. 6218185, an A3 promoter (base numbers 1764-2595 of GenBank registration no. U49854), GFP originating in pEGFP-N1 vector (Clontech) and poly A addition sequence originating in SV40 (base numbers 659-2578 of GenBank registration no. U55762) are inserted into that portion (Nature Biotechnology 18, 81-84, 2000). The expression unit of feline interferon- ω gene was inserted at the XhoI site upstream from the A3 promoter. The expression units of the inserted genes consisted of a sericin-1 gene promoter-feline interferon- ω -bovine growth hormone poly A addition sequence (SEQ. ID No. 1), or a fibroin H chain gene promoter-feline interferon- ω -bovine growth hormone poly A addition sequence (SEQ. ID No. 2). The following provides a detailed description of the method.

Genes were cleaved from the plasmids prepared in Example 2 using the restrictase sites preset in the primers. Namely, insert fragments were cleaved using EcoRI and SalI in the case of sericin-1 gene promoter and fibroin H chain gene promoter, SalI and XbaI in the case of feline interferon- ω , and XbaI and BamHI in the case of bovine growth hormone poly A, followed by electrophoresing with 1 to 1.5% agarose gel and extracting and purifying the fragments in accordance with ordinary methods.

- 30 -

200 ng of sericin-1 gene fragment, 100 ng of feline interferon- ω gene fragment, and 50 ng of bovine growth hormone poly A were mixed and reacted overnight at 16°C by adding an equal volume of DNA Ligation Kit Version 2 (Takara Shuzo). 0.5 μ l of the reaction solution was subjected to PCR using primer 11 (SEQ. ID No. 11) and primer 12 (SEQ. ID No. 12) for 2 minutes of elongation under the same conditions as Example 2. These reaction solutions were electrophoresed with 1% agarose gel, and the amplified, roughly 1.9 kb, DNA fragment (SIB fragment) was extracted and purified in accordance with ordinary methods.

Similarly, 70 ng of fibroin H chain gene promoter fragment, 100 ng of feline interferon- ω gene fragment and 50 ng of bovine growth hormone poly A were mixed and reacted overnight at 16°C by adding an equal volume of DNA Ligation Kit Version 2 (Takara Shuzo). 0.5 μ l of the reaction solution was subjected to PCR using primer 13 (SEQ. ID No. 13) and primer 12 (SEQ. ID No. 12) for 2 minutes of elongation under the same conditions as Example 1. These reaction solutions were electrophoresed with 1% agarose gel, and the amplified roughly 1.5 kb DNA fragment (FIB fragment) was extracted and purified in accordance with ordinary methods.

After digesting these fragments with XhoI, they were ligated to pigA3GFP and subjected to XhoI treatment and dephosphorylation treatment by reacting overnight at 16°C using DNA Ligation Kit Ver. 2 (Takara Shuzo). The plasmid containing the SIB fragment was designated as pigSIB (Fig. 1), and the plasmid containing the FIB fragment was designated as pigFIB (Fig. 2), and these were purified by centrifuging twice using the cesium chloride method and then used in a gene insertion experiment.

Example 4 - Production of Gene Recombinant Silkworms (Fibroin H Chain Gene Promoter)

The aforementioned pigFIB and helper plasmid pHA3PIG (Fig. 3, Nature Biotechnology 18, 81-84, 2000) were adjusted to a concentration of 200 ng/ml each in 0.5 mM phosphate buffer (pH 7.0) and 5 mM KCl, after which 15 to 20 nl were micro-injected into silkworm eggs within 4 hours after being laid.

The larva that hatched from those silkworm eggs were raised, and the resulting adults (G0) were crossed within the same group. By observing the resulting first generation (G1) individuals with fluorescence of green fluorescence protein that had been simultaneously inserted with feline interferon- ω gene, those silkworms that contained the feline interferon- ω gene in their chromosomes were screened. The ratios of the moth groups in which silkworms containing the inserted gene were obtained are shown in Table 1. The silkworm eggs were injected twice, and gene recombinant silkworms were obtained from one moth group by the second injection.

Table 1 Acquisition Status of Gene Recombinant Silkworms (Fibroin Heavy Chain Promoter)

Experiment Group	No. of eggs injected	No. of eggs hatched	No. of adults	No. of sibling mated moths	No. of moth groups positive for feline interferon- ω gene
1	1215	292	220	100	0
2	1326	374	250	123	1

The results of Southern blotting on the gene recombinant silkworms obtained from that moth group are shown in Fig. 4. The method employed for Southern blotting consisted of extracting chromosomal DNA from the G1 generation moths, electrophoresing restrictase-treated samples, and detecting a membrane to which the DNA was transferred by chemiluminescence using the AlkPhos Direct Labeling and Detection System (Amersham-Pharmacia) using a nucleic acid probe specific for feline interferon- ω .

- 32 -

When 11 G1 moths were investigated, feline interferon- ω gene was confirmed to have been inserted into 10 of the silkworms.

Example 5 - Confirmation of Feline Interferon
Production (Fibroin H Chain Promoter)

Since feline interferon- ω has antiviral activity, the presence of feline interferon- ω can be determined according to its activity. Silkworms (G1) of the positive moth group obtained in Example 4 were mated with wild silkworms, and the middle and posterior silk glands excised from fifth instar larva of the resulting generation (G2) were confirmed to be inserted with feline interferon- ω gene. These were then homogenized using 20 mM sodium phosphate buffer (pH 7.0), and the resulting extract was measured using an antiviral activity measuring system that used feline cells. As a result, although antiviral activity was detected for both middle silk glands and posterior silk glands from the silk gland extracts of gene-containing silkworms, activity was not detected from the silk gland extracts of wild silkworms used as the control. Those results are shown in Fig. 5.

Feline interferon- ω is thought to mainly be expressed in posterior silk glands under the control of fibroin H chain promoter. It is believed to subsequently migrate into the middle and anterior silk glands in the same manner as fibroin, and the distribution of physiological activity is considered to coincide with this. On the other hand, there was no antiviral activity detected from silkworms into which the gene was not inserted. This clearly demonstrates that feline interferon- ω protein is expressed while retaining its physiological activity in silkworms into which feline interferon- ω gene has been inserted.

Example 6 - Purification of Feline Interferon

Feline interferon was purified from the extract of

- 33 -

posterior silk glands excised from G2 generation, fifth instar silkworms obtained in Example 5. 1 ml of extract was passed through a HiTrap Blue Sepharose column (Amersham-Pharmacia) followed by washing the column with 10 ml of 20 mM sodium phosphate buffer (pH 7.0). Continuing, the column was eluted with 10 ml of 20 mM sodium phosphate buffer (pH 8.0)-0.5 M NaCl and then with 10 ml of 20 mM sodium phosphate buffer (pH 8.0)-1 M NaCl. The washing fraction, 0.5 M elution fraction and 1 M elution fraction were collected, desalted and concentrated to about 1 ml. The results of determining the antiviral activity and amount of protein of the extract and each purified fraction are shown in Table 2.

Table 2 Purification of Feline Interferon- ω by Blue Sepharose Chromatography

	Antiviral activity (U/ml)	Amt. of protein (mg/ml)	Specific activity (U/mg)
Extracted sample	523	0.37	1401
Blank, washing fraction	23	2.91	8
0.5 M NaCl elution fraction	1494	0.85	1758
1 M NaCl elution fraction	>6270	0.41	>15293

As a result of the purification procedure, antiviral activity, namely feline interferon- ω , could be recovered in the 1 M elution fraction, and its specific activity was roughly 10 times that of the extract.

Example 7 - Production of Recombinant Gene Silkworms (Sericin-1 Promoter)

The aforementioned pigSIB and a helper plasmid were adjusted to a concentration of 200 ng/ml each in 0.5 mM phosphate buffer (pH 7.0) and 5 mM KCl, after which 15 to 20 nl were micro-injected into silkworm eggs within 4 hours after being laid. The larva that hatched from

- 34 -

those silkworm eggs were raised, and the resulting adults (G0) were crossed within the same group. Insertion of feline interferon- ω gene into the chromosomes was investigated by observing the fluorescence of green fluorescence protein from the resulting first-generation (G1) individuals. In two experiments, a gene recombinant vector containing feline interferon- ω gene coupled to sericin promoter was micro-injected into 1218 and 1375 eggs, respectively, and 12 positive moth groups each were able to be obtained (Table 3).

Table 3 Acquisition Status of Gene Recombinant Silkworms (Sericin Promoter)

Experiment Group	No. of eggs injected	No. of eggs hatched	No. of adults	No. of sibling mated moths	No. of moth groups positive for feline interferon- ω gene
1	1218	500	320	158	12
2	1375	540	500	225	12

One silkworm (G1) each that was confirmed to contain the gene was selected from 3 moths groups in the first experiment and 2 moth groups in the second experiment among the resulting positive moth groups, and genomic DNA was extracted from their silk glands. After treating with EcoRI or BglII, Southern blotting analysis was performed on the DNA using feline interferon- ω gene as a probe. Those results are shown in Fig. 6. As a result, feline interferon- ω gene was confirmed to be inserted in the genomes of all silkworms. In addition, the site at which the gene was inserted into the genome was determined to be different depending on the moth group due to differences in the detected signal size.

Next, mRNA expression of feline interferon- ω gene was investigated. Seven G1 silkworms confirmed to contain feline interferon- ω gene by Southern blotting were randomly selected, their mRNA was extracted and the

expression of feline interferon- ω gene mRNA was investigated by RT-PCR. Isogen (Nippon Gene) and Oligodex dT-30 (Roche Diagnostics) were used for mRNA extraction and purification, the Ready-To-Go T-Primed First-Strand Kit (Amersham-Pharmacia) was used for cDNA synthesis, and the procedure was carried out according to the protocol provided with the kit. As a result of carrying out PCR under the same conditions as during the acquisition of feline interferon- ω gene of Example 2, expression of feline interferon- ω gene mRNA was confirmed for all of the silkworms (Fig. 7).

Example 8 - Confirmation of Feline Interferon Production in Middle Silk Glands and Cocoon and Silk Thread

The middle silk glands were excised from three gene recombinant silkworms obtained in Example 7 and one wild silkworm followed by homogenizing using 20 mM sodium phosphate buffer (pH 7.0) and centrifuging to prepare extracts. In addition, one cocoon each from the gene recombinant silkworms and wild silkworm were extracted in the same manner. When these extracts were measured for their antiviral activity, antiviral activity was detected in the middle silk glands of all of the gene recombinant silkworms, but was not detected in the silk glands of the wild silkworm. Moreover, antiviral activity was also detected in the cocoons of the gene recombinant silkworms (Fig. 8).

On the basis of these findings, feline interferon- ω was determined to be expressed in gene recombinant silkworms while retaining its physiological activity, and that activity was determined to remain in the silk thread.

Example 9 - Production of Plasmids for Insertion of Human Interferon- β Gene

Production of plasmids for insertion of human

- 36 -

interferon- β gene was carried out according to the same method as the case of feline interferon- ω gene indicated in Examples 2 through 4.

5 Namely, PCR was carried out using primer 14 (SEQ. ID No. 14) and primer 15 (SEQ. ID No. 15) and using plasmid pORF-hIFN- β encoding human interferon- β gene as a template to obtain a human interferon- β gene fragment. After treating this fragment with restriction endonucleases SalI and XbaI, a plasmid was constructed
10 that contained a gene expression sequence in which fibroin H chain gene promoter was coupled to the 5' terminal or poly A signal originating in bovine growth hormone gene was coupled to the 3' terminal (fibroin H chain promoter-human interferon- β gene-bovine growth
15 hormone gene poly A signal (FhIB): SEQ. ID No. 16, sericin gene promoter-human interferon- β -bovine growth hormone gene poly A signal (ShIB): SEQ. ID No. 17).

The aforementioned FhIB and ShIB sequences for gene expression were then respectively cleaved from these
20 plasmids by treating with XhoI, and coupled to pigA3GFP that had been subjected to dephosphorylation treatment after being cleaved with XhoI. The plasmid containing the FhIB fragment was designated as pigFhIB, while the plasmid that contained the ShIB fragment was designated
25 as pigShIB. These fragments were purified by centrifuging twice according to the cesium chloride method and then used in a gene insertion experiment.

Example 10 - Production of Human Interferon- β Gene Recombinant Silkworms

30 Production of human interferon- β gene recombinant silkworms was carried out according to the same method used to produce feline interferon- ω gene recombinant silkworms indicated in Example 4 by using the gene insertion plasmids produced in Example 9.

- 37 -

Namely, pigFhIB and pigShIB were respectively micro-injected into silkworm eggs together with helper plasmid pHA3PIG, and the resulting adults were crossed followed by screening of the next generation. When each plasmid was injected into 600 eggs, silkworms positive for green fluorescence were obtained in 7 moth groups for silkworms containing pigFhIB and in 5 moth groups for silkworms containing pigShIB, and insertion of the genes into their chromosomes was confirmed by PCR. The silk glands and silk thread were harvested from these silkworms and their extracts were used to measure the physiological activity of human interferon- β in the form of their antiviral activity. The values are shown in the table after having been corrected for total protein concentrations in the samples.

Table 4 Antiviral Activity in Tissue Extracts of Human Interferon- β Gene Recombinant Silkworms

Promoter	Moth group no. - Individual no.	Antiviral activity (units/g protein)		
		Posterior silk glands	Middle silk glands	Silk thread
Fibroin H chain	3-1	599723	82591	Not tested
	3-2	656110	41545	
	11-1	502750	19859	
	11-2	115560	39130	
Sericin	1-1	Not tested	53884	42187
	1-2		648953	101713
	5-1		437291	133288
	5-2		541106	92749
Normal silkworms		Not detected	Not detected	Not detected

Detection limit: Approx. 1000 units/g protein

As a result, since antiviral activity was detected from the posterior and middle silk glands of the silkworms inserted with pigFhIB and antiviral activity was detected from the middle silk glands and silk thread of the silkworms inserted with pigShIB, interferon- β was confirmed to be produced in the silk gland tissue of these silkworms.

- 38 -

Example 11 - Production of Plasmids for Insertion of Feline Granulocyte Colony Stimulating Factor Gene

Production of plasmids for insertion of feline granulocyte colony stimulating factor gene was carried out according to the same method as the case of feline interferon- ω gene indicated in Examples 2 through 4.

Feline granulocyte colony stimulating factor gene was obtained in the form of a feline granulocyte colony stimulating factor gene fragment by carrying out PCR using primer 18 (SEQ. ID No. 18) and primer 19 (SEQ. ID No. 19) from cDNA obtained from CRFK cells stimulated for 24 hours with LPS at 10 μ g/ml according to the report of Yamamoto, et al. (Gene, 274, 263-269, 2001). After treating this fragment with SalI and XbaI, plasmids were constructed that contained a sequence for gene expression in which fibroin H chain gene promoter or sericin gene promoter was coupled to the 5' terminal or poly A signal originating in bovine growth hormone gene was coupled to the 3' terminal (fibroin H chain promoter-feline granulocyte colony stimulating factor gene-bovine growth hormone gene poly A signal (FGB): SEQ. ID No. 20, sericin gene promoter-feline granulocyte colony stimulating factor gene-bovine growth hormone gene poly A signal (SGB): SEQ. ID No. 21).

The aforementioned FGB and SGB sequences for gene expression were then respectively cleaved from these plasmids by treating with XhoI, and coupled to pigA3GFP that had been subjected to dephosphorylation treatment after being cleaved with XhoI. The plasmid containing the FGB fragment was designated as pigFGB, while the plasmid that contained the SGB fragment was designated as pigSGB. These fragments were purified by centrifuging twice according to the cesium chloride method and then used in a gene insertion experiment.

Example 12 - Production of Feline Granulocyte Colony Stimulating Factor Gene Recombinant Silkworms

- 39 -

Production of feline granulocyte colony stimulating factor gene recombinant silkworms was carried out according to the same method used to produce feline interferon- ω gene recombinant silkworms indicated in Example 4 by using the gene insertion plasmids produced in Example 11.

Namely, pigFhIB and pigShIB were respectively micro-injected into silkworm eggs together with helper plasmid pHA3PIG, and the resulting adults were crossed followed by screening of the next generation. When each plasmid was injected into 600 eggs, silkworms positive for green fluorescence were obtained in 3 moth groups for silkworms containing pigFGB and in 7 moth groups for silkworms containing pigSGB, and insertion of the genes into their chromosomes was confirmed by PCR. The silk glands and silk thread were harvested from these silkworms and their extracts were used to measure the physiological activity of feline granulocyte colony stimulating factor in the form of growth promoting activity of NFS-60 cells (ATCC).

Measurement of growth promoting activity was carried out in the manner described below. First, NFS-60 cells were seeded in a 96-well plate in the absence of M-CSF at 2×10^4 cells/well followed 30 minutes later by the addition of 10 μ l of sample. After culturing for an additional 24 hours, cell growth activity was measured using the Cell Counting Kit-8 (Dojindo). The amount of sample that yielded 50% of the maximum growth promoting effect (ED50) was defined as 1 unit/ml, and the physiological activity in the sample was calculated by multiplying by the dilution factor. The values are shown in the table after having been corrected for total protein concentrations in the samples.

- 40 -

Table 5 Growth Promoting Activity in Tissue Extracts of
Feline Granulocyte Colony Stimulating Factor Gene
Recombinant Silkworms

Promoter	Moth group no. - Individual no.	Growth promoting activity (units/g protein)		
		Posterior silk glands	Middle silk glands	Silk thread
Fibroin H chain	9-1	36	Not detected	Not tested
	9-2	412		
	16-1	326	113	
	16-2	4039	226 53	
Sericin	3-1	Not tested	4330	590
	3-2		2277	524
	8-1		3966	846
	8-2		2137	211
Normal silkworms		Not detected	Not detected	Not detected

5 Detection limit: Approx. 20 units/g protein

As a result, as growth promoting activity was
detected from the posterior and middle silk glands of the
silkworms inserted with pigFGB and growth promoting
activity was detected from the middle silk glands and
10 silk thread of the silkworms inserted with pigSGB, feline
granulocyte colony stimulating factor was confirmed to be
produced in the silk gland tissue of these silkworms.

Example 13 - Gene Preparation

A study was conducted on improving production
15 amounts in silk gland tissue and silk thread using feline
interferon- ω as a model of a physiologically active
protein.

The genes used were acquired by PCR by producing
primers for the sequences on both ends using known
20 sequences and using suitable DNA sources for the
templates. Restrictase sites were added to the ends of
the primers for subsequent gene manipulation.

Fibroin H chain promoter (base numbers 62118-62437
of GenBank registration no. AF226688: to be referred to
25 as the P region) was acquired by PCR using two types of
primers consisting of primer 25 (SEQ. ID No. 25) and

- 41 -

primer 26 (SEQ. ID No. 26) and using *Bombyx mori* genomic DNA for the template.

5 Fibroin H chain promoter-fibroin H chain gene first exon-first intron-second exon region (base numbers 62118-63513 of GenBank registration no. AF226688: to be referred to as the HP region) was acquired by using two types of primers consisting of primer 25 (SEQ. ID No. 25) and primer 31 (SEQ. ID No. 31) and using *Bombyx mori* genomic DNA for the template.

10 Fibroin H chain upstream promoter-fibroin H chain gene first exon-first intron region (base numbers 57444-62927 of GenBank registration no. AF226688: to be referred to as the HUP region) was acquired by PCR using two types of primers consisting of primer 33 (SEQ. ID No. 33) and primer 34 (SEQ. ID No. 34) and using *Bombyx mori* genomic DNA for the template.

15 Feline interferon- ω gene (base numbers 9-593 of GenBank registration no. S62636: to be referred to as the IC region) was acquired by PCR using two types of primers consisting of primer 27 (SEQ. ID No. 27) and primer 28 (SEQ. ID No. 28) and using baculovirus rBNV100, which encodes feline interferon- ω gene, for the template. rBNV100 can be produced by, for example, cutting out feline interferon- ω gene from a plasmid extracted from
25 *E. coli* (pFeIFN1) (Patent Microorganism Depository No. 1633), coupling to a silkworm cloning vector (T. Horiuchi, et al., Agric. Biol. Chem., 51, 1573-1580, 1987), and co-transfecting silkworm established cells with the recombinant plasmid produced and silkworm
30 nuclear polyhedrosis virus DNA.

 Fibroin H chain poly A signal region (base numbers 79201-79995 of GenBank registration no. AF226688: to be referred to as the A region) was acquired by PCR using two types of primers consisting of primer 29 (SEQ. ID No. 29) and primer 30 (SEQ. ID No. 30) and using *Bombyx mori* genomic DNA for the template.
35

- 42 -

Fibroin H chain C terminal region gene-fibroin H chain poly A signal region (base numbers 79099-79995 of GenBank registration no. AF226688: to be referred to as the HA region) was acquired by PCR using two types of primers consisting of primer 32 (SEQ. ID No. 32) and primer 30 (SEQ. ID No. 30) and using Bombyx mori genomic DNA for the template.

β -galactosidase (β -gal) gene was acquired by PCR using two types of primers consisting of primer 37 (SEQ. ID No. 37) and primer 38 (SEQ. ID No. 38) and using p β gal-Basic vector (Clontech) for the template.

PCR was carried out in accordance with the accompanying protocol using KODplus (Toyobo). Namely, after adding 100 ng of each template in the case of Bombyx mori genomic DNA or 10 ng in the case of Bombyx mori posterior silk gland cDNA and p β gal-Basic vector, 50 pmol of each primer and 10 μ l of the 10xPCR buffer provided, each reagent was added to a concentration of 1 mM MgCl₂, 0.2 mM dNTPs and 2 units of KODplus followed by bringing to a final volume of 100 μ l. The PCR components were then reacted for 30 cycles using a Perkin-Elmer DNA thermal cycler under DNA denaturation conditions of 94°C for 15 seconds, primer annealing conditions of 55°C for 30 seconds, and elongation conditions of 68°C for 60 to 300 seconds.

These reaction solutions were electrophoresed with 1 to 1.5% agarose gel, and a DNA fragment of roughly 0.3 kbp in the P region, roughly 1.4 kbp in the HP region, roughly 5.5 kbp in the HUP region, roughly 580 bp in the IC region, roughly 0.8 bp in the A region, roughly 0.9 bp in the HA region and roughly 3.2 kbp in the β -gal gene were extracted and prepared in accordance with ordinary methods. After phosphorylating these DNA fragments with polynucleotide kinase (Takara Shuzo), they were ligated to pUC19 vector subjected to dephosphorylation treatment

- 43 -

after being cleaved with HincII by reacting overnight at 16°C using DNA Ligation Kit Ver. 2 (Takara Shuzo). These were then used to transform *E. coli* in accordance with ordinary methods and the resulting transformants were confirmed to be inserted with PCR fragments by performing PCR on the resulting colonies under the same conditions as previously described to prepare plasmids in which the PCR fragments were inserted according to ordinary methods. These plasmids were sequenced to confirm that the resulting fragments consisted of the base sequences of each gene.

Example 14 - Production of Plasmids for Expression of β -Galactosidase

The plasmid retaining β -gal gene prepared in Example 13 was cleaved with SalI and HindIII followed by insertion therein of a roughly 0.3 kbp fragment (P region) cleaved by SalI and HindIII from a plasmid retaining fibroin H chain promoter. Moreover, this was then cleaved with BamHI followed by insertion therein of a roughly 0.8 kbp region (A region) cleaved with BamHI from a plasmid having a fibroin H chain poly A signal region, and purifying the resulting plasmid retaining the β -gal gene using the Qiagen Plasmid Maxi Kit in accordance with the protocol provided. The resulting plasmid was named pPgala, and it was confirmed to be the target plasmid by PCR and sequencing.

Similarly, the plasmid retaining the β -gal gene prepared in Example 13 was cleaved with SalI and HindIII followed by insertion therein of a roughly 1.4 kbp fragment (HP region) cleaved with SalI and HindIII from a plasmid retaining the fibroin H chain promoter-fibroin H chain gene first exon-first intron-second exon region. Moreover, this was then cleaved with BamHI followed by insertion therein of a roughly 0.9 kbp fragment (HA region) cleaved with BamHI from a plasmid retaining a fibroin H chain C terminal region-fibroin H chain poly A

- 44 -

signal region, and purifying the resulting plasmid retaining the β -gal gene using the Qiagen Plasmid Maxi Kit in accordance with the protocol provided. The resulting plasmid was named pHPgalHA, and it was confirmed to be the target plasmid by PCR and sequencing.

Example 15 - Production of Plasmids for Gene Insertion

pigA3GFP (Nature Biotechnology 18, 81-84, 2000) was used for the plasmid for gene insertion. Namely, vector pigA3GFP is a vector in which after removing a region encoding transposase from plasmid p3E1.2 disclosed in US Patent No. 6218185, an A3 promoter (base numbers 1764-2595 of GenBank registration no. U49854), GFP originating in pEGFP-N1 vector (Clontech) and poly A addition sequence originating in SV40 (base numbers 659-2578 of GenBank registration no. U55762) are inserted into that portion. The XhoI site located upstream from the A3 promoter was blunt ended followed by insertion of an expression cassette of feline interferon- ω gene. The constitution of the gene expression cassette used in the present example consisted of fibroin H chain promoter-feline interferon- ω -fibroin H chain C terminal region-fibroin H chain poly A signal region (HP-IC-HA), or fibroin H chain upstream promoter-fibroin H chain gene first exon-first intron-second exon region-feline interferon- ω -fibroin H chain C terminal region-fibroin H chain poly A signal region (HUP-IC-HA), or fibroin H chain promoter-fibroin H chain gene first exon-first intron-second exon region-feline interferon- ω -fibroin H chain poly A signal region (HP-IC-A).

The following indicates the specific method employed.

The P-IC-A construct was produced according to the following procedure. The plasmid retaining feline interferon- ω (IC region) prepared in Example 13 was

cleaved with SalI and HindIII followed by insertion therein of a roughly 0.3 kbp fragment (P region) cleaved with SalI and HindIII from a plasmid retaining fibroin H chain promoter. Moreover, this was cleaved with BamHI followed by insertion therein of a roughly 0.8 kbp fragment (region A) cleaved with BamHI from a plasmid retaining fibroin H chain poly A signal region. This plasmid retaining P, IC and A was cleaved with AscI and the cleaved roughly 1.7 kbp fragment was blunt ended with T4 DNA Polymerase (Takara Shuzo) and coupled to blunt ended and dephosphorylated pigA3GFP XhoI site to produce a construct for gene insertion containing the P-IC-A gene cassette. The procedure is shown in Figs. 9 and 10.

The HP-IC-HA construct was produced in the following manner. The plasmid retaining feline interferon- ω (IC region) prepared in Example 13 was cleaved with SalI and HindIII followed by insertion therein of a roughly 1.4 kbp fragment (HP region) cleaved with SalI and HindIII from a plasmid retaining a fibroin H chain promoter-fibroin H chain gene first exon-first intron-second exon region. Moreover, this was cleaved with BamHI followed by insertion therein of a roughly 0.9 kbp fragment (region HA) cleaved with BamHI from a plasmid retaining fibroin H chain C terminal region-fibroin H chain poly A signal region. This plasmid retaining HP, IC and HA was cleaved with AscI and the cleaved roughly 2.9 kbp fragment was blunt ended with T4 DNA Polymerase (Takara Shuzo) and coupled to blunt ended and dephosphorylated pigA3GFP XhoI site to produce a construct for gene insertion containing the HP-IC-HA gene cassette. The procedure is shown in Figs. 11 and 12.

The HUP-IC-HA construct was produced according to the following procedure. A roughly 2.1 kbp fibroin H chain first intron-second exon region-feline interferon- ω -fibroin H chain C terminal region-fibroin H chain poly A signal region was acquired by PCR using two types of

- 46 -

primers consisting of primer 35 (SEQ. ID No. 35) and primer 36 (SEQ. ID No. 36) and using 1 ng of HP-IC-HA construct for the template. This was then cleaved with XhoI and SphI followed by insertion therein of a roughly 5.5 kbp fragment (HUP region) cleaved with XhoI and SphI from a plasmid retaining fibroin H chain upstream promoter-fibroin H chain gene first exon-first intron. This plasmid retaining HUP, IC and HA was cleaved with AscI and the cleaved roughly 7.6 kbp fragment was blunt ended with T4 DNA Polymerase (Takara Shuzo) and coupled to blunt ended and dephosphorylated pigA3GFP XhoI site to produce a construct for gene insertion containing the HUP-IC-HA gene cassette. The procedure is shown in Figs. 13 and 14.

The HP-IC-A construct was produced according to the procedure described below. The plasmid retaining feline interferon- ω (IC region) prepared in Example 13 was cleaved with SalI and HindIII followed by insertion therein of a roughly 1.4 kbp fragment (HP region) cleaved with SalI and HindIII from a plasmid retaining fibroin H chain promoter-fibroin H chain first exon-first intron-second exon region. Moreover, this was cleaved with BamHI followed by insertion therein of a roughly 0.8 kbp fragment (region A) cleaved with BamHI from a plasmid retaining fibroin H chain poly A signal region. This plasmid retaining HP, IC and A was cleaved with AscI and the cleaved roughly 2.8 kbp fragment was blunt ended with T4 DNA Polymerase (Takara Shuzo) and coupled to blunt ended and dephosphorylated pigA3GFP XhoI site to produce a construct for gene insertion containing the HP-IC-A gene cassette. The procedure is shown in Figs. 15 and 16.

The P-IC-A gene insertion construct, HP-IC-HA gene insertion construct, HUP-IC-HA gene insertion construct and HP-IC-A gene insertion construct were purified using the Qiagen Plasmid Maxi Kit in accordance with the protocol provided.

- 47 -

Example 16 - Expression of β -Galactosidase in
Silkworm Silk Gland

Gold particles having a diameter of 1.6 μ m were washed and sterilized with 100% ethanol and then
5 suspended in sterilized distilled water (60 mg/ml). Incorporation of β -gal gene expression cassettes into silkworm silk glands was carried out using a gene gun. Namely, 50 μ l (0.3 mg) of gold particles, 10 μ g of expression plasmid pPgala or pHPgalHA obtained in Example
10 14, 50 μ l of 2.5 M calcium chloride and 20 μ l of 0.1 M spermidine were successively mixed and after allowing to stand for 30 minutes at room temperature, the mixture was centrifuged to recover the gold particles coated with pHgalC. After washing the resulting metal particles
15 twice with 70% ethanol, they were dispersed in 50 μ l of 100% ethanol. 10 μ l of the suspension of gold particles were placed on a microcarrier and dried. The Model PDS-1000/He (Bio-Rad) was used for the gene gun. The
20 posterior silk glands excised from fifth instar third day silkworm larva were gently washed twice with PBS, placed on a 1% agar plate and sprayed with gold particles coated with DNA at a pressure of 1,100 psi. Following insertion of DNA, the silk glands were transferred to 20 ml of Grace's insect medium and cultured for 2 days at 25°C.
25 After culturing, the culture supernatant and silk gland cells were recovered and confirmed for the expression of β -gal.

Expression was confirmed by Western analysis. The silk gland cells were homogenized in PBS to extract the
30 cell contents. The culture supernatant and cell extract were both adjusted to a total protein concentration of 1.0 mg/ml, and these were then used as samples for SDS-PAGE. After blotting onto a membrane, β -gal protein was detected using the ECL PlusTM Western Blotting Kit

- 48 -

(Amersham-Pharmacia) in accordance with the protocol provided. Namely, the blotted membrane was first blocked overnight at 4°C in blocking solution (5% skim milk, 0.1% Tween20/PBS). The membrane was then washed twice with TPBS (0.1% Tween 20/PBS) and treated for 1 hour at room temperature with anti- β -gal protein antibody (Sigma) diluted 1000-fold with TPBS. The membrane was washed twice with TPBS and additionally washed three times with TPBS for 5 minutes each. After diluting 10000-fold with TPBS, the membrane was treated for 1 hour at room temperature with HRP-labeled anti-rabbit IgG antibody. After washing the membrane twice with TPBS and then three times with TPBS for 5 minutes each, the detection reagents of the ECL PlusTM Western Blotting Detection System (Amersham-Pharmacia) were added (Solution A + Solution B). Luminescence was then exposed onto HyperfilmTMECLTM and developed.

Since β -gal protein was only detected in the silk gland cells and culture supernatant containing pHPgalHA, it was clearly determined that a region other than fibroin H chain promoter, namely fibroin H chain gene first exon-first intron-second exon region, plays an important role in protein synthesis or gene expression within the cells. In addition, secretion outside the cells was also confirmed. Those results are shown in Fig. 17.

Example 17 - Production of Recombinant Gene Silkworms

Each of the gene insertion plasmids described in Example 15 and DNA that produces piggyBac transposase protein (pHA3PIG) were adjusted to a concentration of 200 μ g/ml each in 0.5 mM phosphate buffer (pH 7.0) and 5 mM KCl, after which 3 to 20 nl were micro-injected into silkworm eggs within 4 hours after being laid.

The larva that hatched from those silkworm eggs were raised, and the resulting adults (G0) were crossed within

- 49 -

the same group. By observing the resulting first generation (G1) individuals for fluorescence of jellyfish green fluorescence protein, those silkworms that contained the jellyfish green fluorescence protein gene in their chromosomes were screened. As a result, gene recombinant silkworms were obtained that emitted fluorescent light due to the action of the jellyfish green fluorescence protein.

Example 18 - Expression Analysis of Recombinant Protein in Silk Gland Tissue by Western Analysis

The expression of feline interferon- ω in tissue was investigated by Western analysis after recovering the posterior silk gland tissue from non-transformed silkworms and transformed silkworms (HP-IC-A transformed silkworms, HP-IC-HA transformed silkworms and HUP-IC-HA transformed silkworms). The silkworm posterior silk glands were homogenized in 100 mM sodium phosphate buffer (pH 7.0), and the supernatant was recovered following centrifugation for use as samples. Feline interferon was then detected using the ECL PlusTM Western Blotting Kit (Amersham-Pharmacia) in accordance with the protocol provided. Namely, the blotted membrane was blocked overnight at 4°C in blocking solution (5% skim milk, 0.1% Tween20/PBS). The membrane was then washed twice with TPBS (0.1% Tween 20/PBS) and treated for 1 hour at room temperature with anti-feline interferon antibody diluted 1000-fold with TPBS. The membrane was washed twice with TBPS and additionally washed three times with TPBS for 5 minutes each. After diluting 10000-fold with TPBS, the membrane was treated for 1 hour at room temperature with HRP-labeled anti-rabbit IgG antibody. After washing the membrane twice with TPBS and then three times with TPBS for 5 minutes each, the detection reagents of the ECL PlusTM Western Blotting Detection System (Amersham-Pharmacia) were added (Solution A + Solution B). Luminescence was then exposed onto HyperfilmTMECLTM and developed. As a result, in contrast to signals not being

- 50 -

detected from posterior silk gland tissue of the non-transformed silkworms and P-IC-A construct transformed silkworms, signals were detected from the posterior silk gland tissue of transformed silkworms containing the HP-IC-A construct, HP-IC-HA construct and HUP-IC-HA construct. Based on the results of this experiment, a region other than the fibroin H chain promoter, namely the fibroin H chain gene first exon-first intron-second exon region, was reconfirmed to play an important role in drastically improving protein synthesis or gene expression within silkworm posterior silk gland cells. These results are shown in Fig. 18. The accumulated amount of feline interferon in posterior silk gland tissue in the transformed silkworms containing the HUP-IC-HA gene cassette that contains the 5.5 kbp fibroin H chain 5' terminal region, feline interferon gene and fibroin H chain 3' terminal was higher than the accumulated amount of feline interferon in posterior silk gland tissue in transformed silkworms containing the HP-IC-HA gene cassette that contains the fibroin H chain 5' terminal promoter region, feline interferon gene and fibroin H chain 3' terminal. A gene region that improved the amount of protein produced is thought to be present in a region upstream from the H chain 5' terminal.

Example 19 - Measurement of Recombinant Protein in Silk Thread by Western Analysis

Next, the secretion of exogenous protein, namely feline interferon- ω , in silk thread was investigated.

10 mg each of the cocoons from non-transformed silkworms and transformed silkworms (transformed silkworms containing the HP-IC-A gene, transformed silkworms containing the HP-IC-HA gene, and transformed silkworms containing the HUP-IC-HA gene) were weighed out, and after adding 4 ml of 60% LiSCN and stirring, the cocoons were dissolved by allowing to mix overnight at room temperature. The dissolved cocoons were then diluted 10-fold with 8 M urea, 2% SDS and 5% 2-

- 51 -

mercaptoethanol to prepare samples. The levels of feline interferon in the samples were then detected using the ECL Plus™ Western Blotting Kit (Amersham-Pharmacia) in accordance with the protocol provided. Those results were then measured for signal intensity using a Molecular Imager (Bio-Rad) and the protein contents were measured by comparing with the signal intensities of known concentrations of feline interferon.

As a result, in contrast to signals not being detected from the cocoons of the non-transformed silkworms and transformed silkworms containing HP-IC-A gene, signals were detected from the cocoons of transformed silkworms containing HP-IC-HA gene and transformed silkworms containing HUP-IC-HA gene, thereby confirming that feline interferon protein is secreted into silk thread. In addition, the content was about 0.8 to 2.0% in HP-IC-HA transformed silkworms, and about 1.8 to 5.4% in HUP-IC-HA transformed silkworms. This is equivalent to 0.4 to 2 mg in terms of the weight per silkworm.

On the basis of the results of this experiment, the 3' terminal portion of fibroin H chain gene was clearly demonstrated to play an important role in the secretion of protein synthesized in posterior silk gland cells into silk thread. These results are shown in Fig. 19. The amount of feline interferon produced in transformed silkworms containing the HUP-IC-HA gene cassette that contains a 5.5 kbp fibroin H chain 5' terminal promoter region, feline interferon gene and fibroin H chain 3' terminal was higher than the amount of feline interferon produced in transformed silkworms containing the HP-IC-HA gene cassette that contains fibroin H chain 5' terminal promoter region, feline interferon gene and fibroin H chain 3' terminal. A gene region that improves the amount of protein produced is considered to be present in a region upstream from the H chain 5' terminal.

- 52 -

Example 20 - Measurement of Recombinant Protein in Silk Thread by ELISA

Quantitative determination of feline interferon- ω in silk thread was carried out.

5 10 μ g each of the cocoons from non-transformed silkworms and transformed silkworms (transformed silkworms containing the HP-IC-A gene, transformed silkworms containing the HP-IC-HA gene, and transformed silkworms containing the HUP-IC-HA gene) were weighed
10 out, and after adding 4 ml of 60% LiSCN and stirring, the cocoons were dissolved by allowing them to mix overnight at room temperature. The dissolved cocoons were diluted 8-fold or 16-fold with PBS and applied to a microtiter plate. Known concentrations of feline interferon
15 serially diluted with PBS were used for the standards.

As a result, feline interferon- ω was not detected in silk thread in transformed silkworms containing HP-IC-A gene, but was detected at about 1.1 to 2.2% in transformed silkworms containing HP-IC-HA gene, and at
20 about 1.0 to 4.9% in transformed silkworms containing HUP-IC-HA gene.

INDUSTRIAL APPLICABILITY

25 A cytokine could be recovered in large volume while retaining its physiological activity from the silk glands or silk thread of gene recombinant silkworms obtained by producing a plasmid vector, in which a cytokine gene is coupled to a promoter that functions in silkworm silk glands, and then inserting those genes into silkworm
30 chromosomes. In addition, the resulting cytokine extract has a low level of contaminating proteins, and can be purified easily as compared with methods of the prior art.

35 A large amount of exogenous protein could be produced within silk gland cells, outside silk gland cells and in silk thread by inserting an expression gene

- 53 -

cassette, in which the DNA sequence of the 5' terminal portion and the DNA sequence of the 3' terminal portion of fibroin H chain gene were fused to an exogenous protein gene, into silk gland cells and so forth. The
5 use of this novel technique led to the establishment of a technology for producing easily purified exogenous protein by producing exogenous protein using silk glands without the use of a recombinant baculovirus.

SEQUENCE LISTING

<110>Toray Industries, Inc.

<110>National Institute of Agrobiological Science

<120>Process for production of physiologically active
proteins using recombinant silkworm

<130>

<160>38

<210>1

<211>1910

<212>DNA

<213>Artificial sequence

<220>

<221>CDS

<222>(1071)...(1652)

<221>sig_peptide

<222>(1071)...(1139)

<221>mat_peptide

<222>(1140)...(1652)

<400>1

```

ctcgagggtc agaaacottg ttaaccaata gagccaaata tagttaacac aatagaaatt    60
tatccaaata ttattcgtgt attgtttata gcotttgtca agtottttac aaggcaagat    120
aataagtaat attccgtgat tggacgtaac atttcccgga agatccttag ccgataagtc    180
gaagagccgc atgtggctag agagacgcgg gtttccgacc actggcttag gcgcttattc    240
cgccataata gatgtacgtg ttcacaatta gcaccogaaa ttcgtaatag ctacgagaag    300
tatcgaatat caaaaatcta tatattaata cgtgaagcaa aaactttgta tcccttttta    360
cgaaaattgc gaggacggag gagtatgaaa tttcccacac ttatagagaa tacagagaag    420
aagtgcacaa tgotaatatt tttttaaaat aatgcataaa agatacttta aatcaataaa    480
gaaaacagca cacacactac ataccatgta tttgacgcac acacgcattg atactattta    540
ttgtcaaaact tttgttcttg acgtctgtgt tcaaactgag aatagattaa atattgtttg    600
tctttattaa tatTTTTTaa tagtgtagtc ttggcgaaat ttgtgattat agaagtataa    660
aatacaatca taatagtgtg caaacttaca attcccaatt aattatagtc gaatttcgac    720
tactgcggga cctctagtat taataattct ctttaaaaaa aaacagagca tcaaatactg    780
tcacaaatgt caagcgggtc tcaacgagcc atgaataaat tagaaatcaa ttaataacat    840

```

```

aaaataggca aacaaaataa aaccatttac atagagaacg ttgtttgaac aaaaacaata 900
acttgtatac attgtttgca caaatgtttg aaccgaaaat ttattactct ctacgtaagc 960
ttgatcaaac ttctgtttcg tataaaacgc gttggcccaa ccactttggc atagtgtct 1020
tatcatcggtg tctctaagga tcaagcgatc caaagaccgc caacgtcgac atg gcg 1076
Met Ala
1
ctg ccc tct tcc ttc ttg gtg gcc ctg gtg gcg ctg ggc tgc aac tcc 1124
Leu Pro Ser Ser Phe Leu Val Ala Leu Val Ala Leu Gly Cys Asn Ser
5 10 15
gtc tgc gtg ctg ggc tgt gac ctg cct cag acc cac ggc ctg ctg aac 1172
Val Cys Val Leu Gly Cys Asp Leu Pro Gln Thr His Gly Leu Leu Asn
20 25 30
agg agg gcc ttg acg ctc ctg gga caa atg agg aga ctc cct gcc agc 1220
Arg Arg Ala Leu Thr Leu Leu Gly Gln Met Arg Arg Leu Pro Ala Ser
35 40 45 50
tcc tgt cag aag gac aga aat gac ttc gcc ttc ccc cag gac gtg ttc 1268
Ser Cys Gln Lys Asp Arg Asn Asp Phe Ala Phe Pro Gln Asp Val Phe
55 60 65
ggt gga gac cag tcc cac aag gcc caa gcc ctc tcg gtg gtg cac gtg 1316
Gly Gly Asp Gln Ser His Lys Ala Gln Ala Leu Ser Val Val His Val
70 75 80
acg aac cag aag atc ttc cac ttc ttc tgc aca gag gcg tcc tcg tct 1364
Thr Asn Gln Lys Ile Phe His Phe Phe Cys Thr Glu Ala Ser Ser Ser
85 90 95
gct gct tgg aac acc acc ctc ctg gag gaa ttt tgc acg gga ctt gat 1412
Ala Ala Trp Asn Thr Thr Leu Leu Glu Glu Phe Cys Thr Gly Leu Asp
100 105 110
cgg cag ctg acc cgc ctg gaa gcc tgt gtc ctg cag gag gtg gag gag 1460
Arg Gln Leu Thr Arg Leu Glu Ala Cys Val Leu Gln Glu Val Glu Glu
115 120 125 130
gga gag gct ccc ctg acg aac gag gac att cat ccc gag gac tcc atc 1508
Gly Glu Ala Pro Leu Thr Asn Glu Asp Ile His Pro Glu Asp Ser Ile
135 140 145

```

```

ctg agg aac tac ttc caa aga ctc tcc ctc tac ctg caa gag aag aaa      1556
Leu Arg Asn Tyr Phe Gln Arg Leu Ser Leu Tyr Leu Gln Glu Lys Lys
      150                      155                      160
tac agc cct tgt gcc tgg gag atc gtc aga gca gaa atc atg aga tcc      1604
Tyr Ser Pro Cys Ala Trp Glu Ile Val Arg Ala Glu Ile Met Arg Ser
      165                      170                      175
ttg tat tat tca tca aca gcc ttg cag aaa aga tta agg ago gag aaa      1652
Leu Tyr Tyr Ser Ser Thr Ala Leu Gln Lys Arg Leu Arg Ser Glu Lys
      180                      185                      190
tga tctagaccgc tgatcagcct cgactgtgcc ttctagttgc cagccatctg      1705
ttgtttgccc ctcccccggtg ccttccttga ccctggaagg tgccactccc actgtccttt      1765
cctaataaaa tgaggaaatt gcatogcatt gtctgagtag gtgtcattot attctggggg      1825
gtgggggtggg gcaggacagc aagggggagg attgggaaga caatagcagg catgctgggg      1885
atgcgggtggg ctctatggcc tcgag      1910
<210>2
<211>1172
<212>DNA
<213>Artificial sequence
<220>
<221>CDS
<222>(333)...(914)
<221>sig_peptide
<222>(333)...(401)
<221>mat_peptide
<222>(402)...(914)
<400>2
ctcgagggga gaaagcatga agtaagttct ttaaataatta caaaaaaatt gaacgatatt      60
ataaaattct taaaaatatt aaaagtaaga acaataagat caattaaatc ataattaatc      120
acattgttca tgatcacaat ttaatttact tcatacgttg tattgttatg ttaaataaaa      180
agattaattt ctatgtaatt gtatctgtac aatacaatgt gtagatgttt attctatcga      240
aagtaaatac gtcaaaaactc gaaaattttc agtataaaaa ggttcaactt ttcaaatca      300
gcatacgttc ggttccaact ctcaaggtcg ac atg gcg ctg ccc tot tcc ttc      353

```

Met Ala Leu Pro Ser Ser Phe																
1					5											
ttg	gtg	gcc	ctg	gtg	gcg	ctg	ggc	tgc	aac	tcc	gtc	tgc	gtg	ctg	ggc	401
Leu	Val	Ala	Leu	Val	Ala	Leu	Gly	Cys	Asn	Ser	Val	Cys	Val	Leu	Gly	
10					15					20						
tgt	gac	ctg	cct	cag	acc	cac	ggc	ctg	ctg	aac	agg	agg	gcc	ttg	acg	449
Cys	Asp	Leu	Pro	Gln	Thr	His	Gly	Leu	Leu	Asn	Arg	Arg	Ala	Leu	Thr	
25					30					35						
ctc	ctg	gga	caa	atg	agg	aga	ctc	cct	gcc	agc	tcc	tgt	cag	aag	gac	497
Leu	Leu	Gly	Gln	Met	Arg	Arg	Leu	Pro	Ala	Ser	Ser	Cys	Gln	Lys	Asp	
40					45					50					55	
aga	aat	gac	ttc	gcc	ttc	ccc	cag	gac	gtg	ttc	ggt	gga	gac	cag	tcc	545
Arg	Asn	Asp	Phe	Ala	Phe	Pro	Gln	Asp	Val	Phe	Gly	Gly	Asp	Gln	Ser	
60					65					70						
cac	aag	gcc	caa	gcc	ctc	tcg	gtg	gtg	cac	gtg	acg	aac	cag	aag	atc	593
His	Lys	Ala	Gln	Ala	Leu	Ser	Val	Val	His	Val	Thr	Asn	Gln	Lys	Ile	
75					80					85						
ttc	cac	ttc	ttc	tgc	aca	gag	gcg	tcc	tcg	tct	gct	gct	tgg	aac	acc	641
Phe	His	Phe	Phe	Cys	Thr	Glu	Ala	Ser	Ser	Ser	Ala	Ala	Trp	Asn	Thr	
90					95					100						
acc	ctc	ctg	gag	gaa	ttt	tgc	acg	gga	ott	gat	cgg	cag	ctg	acc	cgc	689
Thr	Leu	Leu	Glu	Glu	Phe	Cys	Thr	Gly	Leu	Asp	Arg	Gln	Leu	Thr	Arg	
105					110					115						
ctg	gaa	gcc	tgt	gtc	ctg	cag	gag	gtg	gag	gag	gga	gag	gct	ccc	ctg	737
Leu	Glu	Ala	Cys	Val	Leu	Gln	Glu	Val	Glu	Glu	Gly	Glu	Ala	Pro	Leu	
120					125					130					135	
acg	aac	gag	gac	att	cat	ccc	gag	gac	tcc	atc	ctg	agg	aac	tac	ttc	785
Thr	Asn	Glu	Asp	Ile	His	Pro	Glu	Asp	Ser	Ile	Leu	Arg	Asn	Tyr	Phe	
140					145					150						
caa	aga	ctc	tcc	ctc	tac	ctg	caa	gag	aag	aaa	tac	agc	cct	tgt	gcc	833
Gln	Arg	Leu	Ser	Leu	Tyr	Leu	Gln	Glu	Lys	Lys	Tyr	Ser	Pro	Cys	Ala	
155					160					165						
tgg	gag	atc	gtc	aga	gca	gaa	atc	atg	aga	tcc	ttg	tat	tat	tca	tca	881

Trp Glu Ile Val Arg Ala Glu Ile Met Arg Ser Leu Tyr Tyr Ser Ser
 170 175 180
 aca gcc ttg cag aaa aga tta agg agc gag aaa tga tctagaccgc 927
 Thr Ala Leu Gln Lys Arg Leu Arg Ser Glu Lys
 185 190
 tgatcagcct cgactgtgcc ttctagttgc cagccatctg ttgtttgccc ctcccccggtg 987
 ccttccttga ccctggaagg tgccactccc actgtccttt cctaataaaa tgaggaaatt 1047
 gcatcgcatg gtotgagtag gtgtcattct attctggggg gtgggggtggg gcaggacagc 1107
 aagggggagg attgggaaga caatagcagg catgctgggg atgcgggtggg ctctatggcc 1167
 tcgag 1172
 <210>3
 <211>31
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>3
 acgcgtcgac atgggcgctg ccctcttcc t 31
 <210>4
 <211>30
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>4
 ctagtctaga tcatttctcg ctcccttaac 30
 <210>5
 <211>29
 <212>DNA
 <223>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>5

cgggaattcg gtcagaaacc ttgttaacc

29

<210>6

<211>30

<212>DNA

<213>Artificial sequence

<220>

<223>Synthesized oligonucleotide

<400>6

acgcgtcgac gttggcggtc ttggatcgc

30

<210>7

<211>29

<212>DNA

<213>Artificial sequence

<220>

<223>Synthesized oligonucleotide

<400>7

cgggaattcg ggagaaagca tgaagtaag

29

<210>8

<211>30

<212>DNA

<213>Artificial sequence

<220>

<223>Synthesized oligonucleotide

<400>8

acgcgtcgac cttgagagtt ggaaccgaac

30

<210>9

<211>30

<212>DNA

<213>Artificial sequence

<220>

<223>Synthesized oligonucleotide

<400>9

ctagtctaga ccgctgatca gcctcgactg

30

<210>10
 <211>30
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>10
 cgcggatccg ccatagagcc caccgcatcc 30
 <210>11
 <211>34
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>11
 gacctcgagg gtcagaaacc ttgttaacca atag 34
 <210>12
 <211>30
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>12
 gacctcgagg ccatagagcc caccgcatcc 30
 <210>13
 <211>29
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>13
 gacctcgagg ggagaaagca tgaagtaag 29
 <210>14

<211>31
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>14
 acgcgtogac atgaccaaca agtgtctcct c 31
 <210>15
 <211>30
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>15
 ctagtctaga tcagtttcgg aggtaacctg 30
 <210>16
 <211>1151
 <212>DNA
 <213>Artificial sequence
 <220>
 <221>CDS
 <222>(333)...(893)
 <221>sig_peptide
 <222>(333)...(395)
 <221>mat_peptide
 <222>(396)...(893)
 <400>16
 ctcgagggga gaaagcatga agtaagttct ttaaattatta caaaaaaatt gaacgatatt 60
 ataaaaattct taaaaatatt aaaagtaaga acaataagat caattaaatc ataattaatc 120
 acattgttca tgatcacaaat ttaatttact tcatacgttg tattgttatg ttaaataaaa 180
 agattaattht ctatgtaatt gtatctgtac aatacaatgt gtagatgttt attctatoga 240
 aagtaaatac gtcaaaaactc gaaaattttc agtataaaaa ggttcaactt tttcaaatca 300
 gcatcagttc gggttcaact ctcaaggctc ac 332

atg acc aac aag tgt ctc ctc caa att gct ctc ctg ttg tgc ttc toc	380
Met Thr Asn Lys Cys Leu Leu Gln Ile Ala Leu Leu Leu Cys Phe Ser	
1 5 10 15	
act aca gct ctt tcc atg agc tac aac ttg ctt gga ttc cta caa aga	428
Thr Thr Ala Leu Ser Met Ser Tyr Asn Leu Leu Gly Phe Leu Gln Arg	
20 25 30	
agc agc aat ttt cag tgt cag aag ctc ctg tgg caa ttg aat ggg agg	476
Ser Ser Asn Phe Gln Cys Gln Lys Leu Leu Trp Gln Leu Asn Gly Arg	
35 40 45	
ctt gaa tac tgc ctc aag gac agg atg aac ttt gac atc cct gag gag	524
Leu Glu Tyr Cys Leu Lys Asp Arg Met Asn Phe Asp Ile Pro Glu Glu	
50 55 60	
att aag cag ctg cag cag ttc cag aag gag gac gcc gca ttg acc atc	572
Ile Lys Gln Leu Gln Gln Phe Gln Lys Glu Asp Ala Ala Leu Thr Ile	
65 70 75 80	
tat gag atg ctc cag aac atc ttt gct att ttc aga caa gat tca tot	620
Tyr Glu Met Leu Gln Asn Ile Phe Ala Ile Phe Arg Gln Asp Ser Ser	
85 90 95	
agc act ggc tgg aat gag act att gtt gag aac ctc ctg gct aat gtc	668
Ser Thr Gly Trp Asn Glu Thr Ile Val Glu Asn Leu Leu Ala Asn Val	
100 105 110	
tat cat cag ata aac cat ctg aag aca gtc ctg gaa gaa aaa ctg gag	716
Tyr His Gln Ile Asn His Leu Lys Thr Val Leu Glu Glu Lys Leu Glu	
115 120 125	
aaa gaa gat ttc acc agg gga aaa ctc atg agc agt ctg cac ctg aaa	764
Lys Glu Asp Phe Thr Arg Gly Lys Leu Met Ser Ser Leu His Leu Lys	
130 135 140	
aga tat tat ggg agg att ctg cat tac ctg aag gcc aag gag tac agt	812
Arg Tyr Tyr Gly Arg Ile Leu His Tyr Leu Lys Ala Lys Glu Tyr Ser	
145 150 155 160	
cac tgt gcc tgg acc ata gtc aga gtg gaa atc cta agg aac ttt tac	860
His Cys Ala Trp Thr Ile Val Arg Val Glu Ile Leu Arg Asn Phe Tyr	
165 170 175	

ttc att aac aga ctt aca ggt tac ctc cga aac tga tctagaccgc 906
 Phe Ile Asn Arg Leu Thr Gly Tyr Leu Arg Asn

180

185

tgatcagcct cgactgtgoc ttctagttgc cagccatctg ttgtttgccc ctccccogtg 960
 ccttccttga ccctggaagg tgcoactccc actgtccttt cctaataaaa tgaggaaatt 1026
 gcacgcatt gtctgagtag gtgtcattct attctggggg gtgggggtggg gcaggacagc 1086
 aagggggagg attggaaga caatagcagg catgctgggg atgcggtggg ctctatggcc 1146
 togag 1151

<210>17

<211>1849

<212>DNA

<213>Artificial sequence

<220>

<221>CDS

<222>(1071)...(1631)

<221>sig_peptide

<222>(1071)...(1133)

<221>mat_peptide

<222>(1134)...(1631)

<400>17

ctogagggtc agaaacottg ttaaccaata gagccaaata tagttaacac aatagaaatt 60
 tatccaaata ttattogtgt attgtttata gcctttgtca agtcttttac aaggcaagat 120
 aataagtaat attcogtgat tggacgtaac atttccogga agatccttag ccgataagtc 180
 gaagagccgc atgtggctag agagacgagg gtttccgacc actggcttag gcgcttattc 240
 cgccataata gatgtaogtg ttcacaatta gcaaccogaaa ttogtaatag ctacgagaag 300
 tatogaatat caaaaatcta tatattaata cgtgaagcaa aaactttgta tcccttttta 360
 cgaaaattgc gaggaaggag gagtatgaaa tttccacac ttatagagaa tacagagaag 420
 aagtgcacaa tgctaataatt tttttaaaat aatgcataaa agatacttta aatcaataaa 480
 gaaaacagca cacacactac ataccatgta tttagcgcae acacgcatgt atactattta 540
 ttgtcaaaact tttgttcttg acgtctgtgt tcaaaactgag aatagattaa atattgtttg 600
 totttattaa tattttttta tagtgtagtc ttggcgaaat ttgtgattat agaagtataa 660
 aatacaatca taatagtgtg caaacttaca attcccaatt aattatagtc gaatttcgac 720
 tactgcggga cctctagtat taataattct ctttaaaaaa aaacagagca tcaaatactg 780

```

tcacaaatgt caagcgggtc tcaacgagcc atgaataaat tagaaatcaa ttaataacat 840
aaaaataggca aacaaaataa aaccatttac atagagaacg tttgttgaaac aaaaacaata 900
acttgtatac attgtttgca caaatgtttg aacogaaaat ttattactct ctacgtaagc 960
ttgatcaaac ttogttttog tataaaacgc gttggcccaa ccactttggc atagtcgtct 1020
tatcatoggg totctaagga tcaagcgatc caaagaccgc caacgtcgac 1070
atg acc aac aag tgt ctc ctc caa att gct ctc ctg ttg tgc ttc tcc 1118
Met Thr Asn Lys Cys Leu Leu Gln Ile Ala Leu Leu Leu Cys Phe Ser
1 5 10 15
act aca gct ctt tcc atg agc tac aac ttg ctt gga ttc cta caa aga 1166
Thr Thr Ala Leu Ser Met Ser Tyr Asn Leu Leu Gly Phe Leu Gln Arg
20 25 30
agc agc aat ttt cag tgt cag aag ctc ctg tgg caa ttg aat ggg agg 1214
Ser Ser Asn Phe Gln Cys Gln Lys Leu Leu Trp Gln Leu Asn Gly Arg
35 40 45
ctt gaa tac tgc ctc aag gac agg atg aac ttt gac atc cct gag gag 1262
Leu Glu Tyr Cys Leu Lys Asp Arg Met Asn Phe Asp Ile Pro Glu Glu
50 55 60
att aag cag ctg cag cag ttc cag aag gag gac gcc gca ttg acc atc 1310
Ile Lys Gln Leu Gln Gln Phe Gln Lys Glu Asp Ala Ala Leu Thr Ile
65 70 75 80
tat gag atg ctc cag aac atc ttt gct att ttc aga caa gat tca tot 1358
Tyr Glu Met Leu Gln Asn Ile Phe Ala Ile Phe Arg Gln Asp Ser Ser
85 90 95
agc act ggc tgg aat gag act att gtt gag aac ctc ctg gct aat gtc 1406
Ser Thr Gly Trp Asn Glu Thr Ile Val Glu Asn Leu Leu Ala Asn Val
100 105 110
tat cat cag ata aac cat ctg aag aca gtc ctg gaa gaa aaa ctg gag 1454
Tyr His Gln Ile Asn His Leu Lys Thr Val Leu Glu Glu Lys Leu Glu
115 120 125
aaa gaa gat ttc acc agg gga aaa ctc atg agc agt ctg cac ctg aaa 1502
Lys Glu Asp Phe Thr Arg Gly Lys Leu Met Ser Ser Leu His Leu Lys
130 135 140
aga tat tat ggg agg att ctg cat tac ctg aag gcc aag gag tac agt 1550

```

Arg Tyr Tyr Gly Arg Ile Leu His Tyr Leu Lys Ala Lys Glu Tyr Ser
 145 150 155 160
 cac tgt gcc tgg acc ata gtc aga gtg gaa atc cta agg aac ttt tac 1598
 His Cys Ala Trp Thr Ile Val Arg Val Glu Ile Leu Arg Asn Phe Tyr
 165 170 175
 ttc att aac aga ctt aca ggt tac ctc cga aac tga cagccatctg 1644
 Phe Ile Asn Arg Leu Thr Gly Tyr Leu Arg Asn
 180 185
 ttgtttgcc ctcctccgtg ccttccttga ccctggaagg tgccactccc actgtccttt 1704
 cctaataaaa tgaggaaatt gcatcgcatg gtctgagtag gtgtcattct attctggggg 1764
 gtggggtggg gcaggacagc aagggggagg attgggaaga caatagcagg catgctgggg 1824
 atgcggtggg ctctatggcc tcgag 1849
 <210>18
 <211>28
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>18
 acgcgctgac atgaagctga ccgcoctg 28
 <210>19
 <211>28
 <212>DNA
 <213>Artificial sequence
 <220>
 <223>Synthesized oligonucleotide
 <400>19
 ctagtctaga tcagggcttg gtgaagtg 28
 <210>20
 <211>1175
 <212>DNA
 <213>Artificial sequence
 <220>

<221>CDS

<222>(333)...(917)

<221>sig_peptide

<222>(333)...(401)

<221>mat_peptide

<222>(402)...(917)

<400>20

```

ctcgaggga gaaagcatga agtaagttct ttaaataatta caaaaaaatt gaacgatatt      60
ataaaattct ttaaaatatt aaaagtaaga acaataagat caattaaatc ataattaatc      120
acattgttca tgatcacaat ttaatttact tcatacgttg tattgttatg ttaaataaaa      180
agattaatatt ctatgtaatt gtatctgtac aatacaatgt gtagatgttt attctatcga      240
aagtaaatac gtcaaaaactc gaaaattttc agtataaaaa ggttcaactt tttcaaatca      300
gcatcagttc ggttccaact ctcaagggtcg ac                                    332
atg aag ctg acc gcc ctg cag ctg ctg ctg tgg cac agc gca ctc tgg      380
Met Lys Leu Thr Ala Leu Gln Leu Leu Leu Trp His Ser Ala Leu Trp
      1              5              10              15
atg gtg caa gaa gcc acc ccc ttg ggc cct acc agc tcc ctg ccc cag      428
Met Val Gln Glu Ala Thr Pro Leu Gly Pro Thr Ser Ser Leu Pro Gln
      20              25              30
agc ttc ctg ctc aag tgc tta gaa caa gtg agg aag gtc cag gct gat      476
Ser Phe Leu Leu Lys Cys Leu Glu Gln Val Arg Lys Val Gln Ala Asp
      35              40              45
ggc aca gcg ctg cag gag agg ctg tgc gcc gcc cac aag ctg tgc cac      524
Gly Thr Ala Leu Gln Glu Arg Leu Cys Ala Ala His Lys Leu Cys His
      50              55              60
cct gag gag ctg gtg ctg ctt ggg cac gct ctg ggc atc ccc cag gct      572
Pro Glu Glu Leu Val Leu Leu Gly His Ala Leu Gly Ile Pro Gln Ala
      65              70              75              80
ccc ctg agc agc tgc tcc agc cag gcc ctg cag ctg acg ggc tgc ctg      620
Pro Leu Ser Ser Cys Ser Ser Gln Ala Leu Gln Leu Thr Gly Cys Leu
      85              90              95
cgt caa ctc cac agt ggc ctc ttc ctc tac cag ggc ctc ctg cag gcc      668

```

```

Arg Gln Leu His Ser Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala
      100              105              110
ctg gca ggg ata tcc ccc gag tta gcc ccc acc ctg gac atg ctg cag      716
Leu Ala Gly Ile Ser Pro Glu Leu Ala Pro Thr Leu Asp Met Leu Gln
      115              120              125
ctg gac atc acc gac ttt gct atc aac atc tgg cag cag atg gaa gac      764
Leu Asp Ile Thr Asp Phe Ala Ile Asn Ile Trp Gln Gln Met Glu Asp
      130              135              140
gtg ggg atg gcc cct gca gtg ccg ccc acc cag ggc acc atg cca acc      812
Val Gly Met Ala Pro Ala Val Pro Pro Thr Gln Gly Thr Met Pro Thr
      145              150              155              160
ttc acc tcg gcc ttc cag cgc cgg gca gga ggc acc ctg gtt gcc tcc      860
Phe Thr Ser Ala Phe Gln Arg Arg Ala Gly Gly Thr Leu Val Ala Ser
      165              170              175
aac ctg cag agc ttc ctg gag gtg gca tac cgt gct ctg cgc cac ttc      908
Asn Leu Gln Ser Phe Leu Glu Val Ala Tyr Arg Ala Leu Arg His Phe
      180              185              190
acc aag ccc tga tctagaccgc tgatcagcct cgactgtgcc ttctagttgc      960
Thr Lys Pro
      195
cagccatctg ttgtttgcc ctcctccgtg ccttccttga ccttgaagg tgccactccc 1020
actgtccttt cctaataaaa tgaggaaatt gcatcgcat gtctgagtag gtgtcattct 1080
attctggggg gtgggggtgg gcaggacagc aagggggagg attgggaaga caatagcagg 1140
catgtggggg atgoggtggg ctctatggcc togag      1175
<210>21
<211>1873
<212>DNA
<213>Artificial sequence
<220>
<221>CDS
<222>(1071)...(1655)
<221>sig_peptide
<222>(1071)...(1139)

```

<221>mat_peptide

<222>(1140)...(1655)

<400>21

```

ctcgagggtc agaaaccttg ttaaccaata gagccaaata tagttaacac aatagaaatt      60
tatccaaata ttattcgtgt attgtttata gcctttgtca agtcttttac aaggcaagat    120
aataagtaat attcogtgat tggacgtaac atttcccgga agatccttag ccgataagtc    180
gaagagccgc atgtggctag agagacgcgg gtttccgacc actggccttag gcgcttattc    240
cgccataata gatgtacgtg ttcacaatta gcaccogaaa ttogtaatag ctacgagaag    300
tatcgaatat caaaaatota tatattaata ogtgaagcaa aaactttgta tcccttttta    360
cgaaaattgc gaggacggag gagtatgaaa tttcccacac ttatagagaa tacagagaag    420
aagtgcacaa tgctaattatt tttttaaaat aatgcataaa agatacttta aatcaataaa    480
gaaaacagca cacacactac ataccatgta ttgacgcac acacgcattgt atactattta    540
ttgtcaaact tttgttcttg acgtctgtgt tcaaactgag aatagattaa atattgtttg    600
tctttattaa tattttttta tagtgtagtc ttggcgaaat ttgtgattat agaagtataa    660
aatacaatca taatagtgtg caaacttaca attcccaatt aattatagtc gaatttcgac    720
tactgoggga cotctagtat taataattct ctttaaaaaa aaacagagca tcaaatactg    780
tcacaaatgt caagcgggtc tcaacgagcc atgaataaat tagaaatcaa ttaataacat    840
aaaataggca aacaaaataa aaccattttac atagagaacg ttgttgaac aaaaacaata    900
acttgtatac attgtttgca caaatgtttg aaccgaaaat ttattactct ctacgtaagc    960
ttgatcaaac ttcgttttcg tataaaacgc gttggcccaa ccactttggc atagtogtct   1020
tatcatcggg totctaagga tcaagcgatc caaagacgcg caacgtcgac               1070
atg aag ctg acc gcc ctg cag ctg ctg ctg tgg cac agc gca ctc tgg      1118
Met Lys Leu Thr Ala Leu Gln Leu Leu Leu Trp His Ser Ala Leu Trp
      1              5              10              15
atg gtg caa gaa gcc acc ccc ttg ggc cct acc agc tcc ctg ccc cag      1166
Met Val Gln Glu Ala Thr Pro Leu Gly Pro Thr Ser Ser Leu Pro Gln
      20              25              30
agc ttc ctg ctc aag tgc tta gaa caa gtg agg aag gtc cag gct gat      1214
Ser Phe Leu Leu Lys Cys Leu Glu Gln Val Arg Lys Val Gln Ala Asp
      35              40              45
ggc aca gcg ctg cag gag agg ctg tgc gcc gcc cac aag ctg tgc cac      1262
Gly Thr Ala Leu Gln Glu Arg Leu Cys Ala Ala His Lys Leu Cys His
      50              55              60

```

```

cct gag gag ctg gtg ctg ctt ggg cac gct ctg ggc atc ccc cag gct      1310
Pro Glu Glu Leu Val Leu Leu Gly His Ala Leu Gly Ile Pro Gln Ala
  65              70              75              80
ccc ctg agc agc tgc toc agc cag gcc ctg cag ctg acg ggc tgc ctg      1358
Pro Leu Ser Ser Cys Ser Ser Gln Ala Leu Gln Leu Thr Gly Cys Leu
              85              90              95
cgt caa ctc cac agt ggc ctc ttc ctc tac cag ggc ctc ctg cag gcc      1406
Arg Gln Leu His Ser Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala
              100              105              110
ctg gca ggg ata tcc ccc gag tta gcc ccc acc ctg gac atg ctg cag      1454
Leu Ala Gly Ile Ser Pro Glu Leu Ala Pro Thr Leu Asp Met Leu Gln
              115              120              125
ctg gac atc acc gac ttt gct atc aac atc tgg cag cag atg gaa gac      1502
Leu Asp Ile Thr Asp Phe Ala Ile Asn Ile Trp Gln Gln Met Glu Asp
              130              135              140
gtg ggg atg gcc cct gca gtg cgg ccc acc cag ggc acc atg cca acc      1550
Val Gly Met Ala Pro Ala Val Pro Pro Thr Gln Gly Thr Met Pro Thr
              145              150              155              160
ttc acc tcg gcc ttc cag cgc cgg gca gga ggc acc ctg gtt gcc toc      1598
Phe Thr Ser Ala Phe Gln Arg Arg Ala Gly Gly Thr Leu Val Ala Ser
              165              170              175
aac ctg cag agc ttc ctg gag gtg gca tac cgt gct ctg cgc cac ttc      1646
Asn Leu Gln Ser Phe Leu Glu Val Ala Tyr Arg Ala Leu Arg His Phe
              180              185              190
acc aag ccc tga cagccatctg ttgtttgcc ctcocccgtg ccttccttga      1698
Thr Lys Pro
              195
ccctggaagg tgccactccc actgtccttt cctaataaaa tgaggaaatt gcatcgcatt      1758
gtctgagtag gtgtcattct attctggggg gtggggtggg gcaggacagc aagggggagg      1818
attggaaga caatagcagg catgctgggg atgcggtggg ctctatggcc tcgag      1873
<210>22
<211>1396
<212>DNA

```


<213>Bombyx mori

<400>22

```

gggagaaagc atgaagtaag ttctttaaat attacaaaaa aattgaacga tattataaaa      60
ttctttaaaa tattaagaat aagaacaata agatcaatta aatcataatt aatcacattg      120
ttcatgatca caatttaatt tacttcatac gttgtattgt tatgttaaatt aaaaagatta      180
atttctatgt aattgtatct gtacaataca atgtgtagat gtttattota tcgaaagtaa      240
atacgtaaaa actogaaaaat tttcagtata aaaagggtca actttttcaa atcagcatca      300
gttcgggttc aactctcaag atgagagtcg aaacctttgt gatcttgtgc tgcgctctgc      360
aggtgagtta attattttac tattatttca gaagggtggc agacgatata acgggccacc      420
tgataataag tggtcgcca aacgcacaga tatcgtaaat tgtgccattt gatttgtcac      480
gccccggggg gctacggaat aaactacatt tatttattta aaaaatgaac cttagattat      540
gtaacttgtg atttatttgc gtcaaaagta ggcaagatga atctatgtaa atacctgggc      600
agacttgcaa tatctattt caccggtaaa tcagcattgc aatatgcaat gcatattcaa      660
caatatgtaa aacaattcgt aaagcatcat tagaaaaatg acgaaagaaa ttgcataaaa      720
ttataacogc attattaatt tattatgata tctattaaca attgctattg cctttttttc      780
gcaattata atcattttca taacctcgag gtagcattct gttacatttt aatacattgg      840
tatgtgatta taacacgagc tgcccaactga gtttctcgcc agatcttctc agtgggtcgc      900
gttaocgata acgtgataga ttctatgaag cactgctctt gttagggtta gtgttagcaa      960
attctttcag gttgagtctg agagctcacc taccatcggg agcgtagctg gaataggcta     1020
ccagctaata ggtagggaaa acaaagctcg aaacaagctc aagtaataac aacataatgt     1080
gaccataaaa tctogtggtg tatgagatac aattatgtac tttcccacaa atgtttacat     1140
aattagaatg ttgttcaact tgcctaacgc ccagctaga acattcaatt attactatta     1200
ccactactaa ggcagtatgt cctaactcgt tccagatcag cgctaacttc gattgaatgt     1260
gcgaaattta tagctcaata ttttagcact tatcgtattg atttaagaaa aaattgttaa     1320
cattttgttt cagtatgtcg cttatacaaa tgcaaacatc aatgatattg atgaggacta     1380
ttttgggagt gatgtc                                     1396

```

<210>23

<211>6070

<212>DNA

<213>Bombyx mori

<400>23

```

tcaagacatc cttgattaag gcagctgccg atattgacat ggacctcgtt cgtgctgcga      60
tagacgactg gccgcgcaga ttgaaggcct gtattcaaaa tcacggaggt cattttgaat     120

```

```

aaactttagt gtcataagaa tctatgtttt gttaagttca ttttggtata tgaatggtta 180
cataatgaat aaacttggtt caattatttt acattaaaca tgtgacagaa tttatgacct 240
gactaggtag gtacaaacag cctttttgat attagaaaac taagtaaaat agcctacggt 300
cacatctctt tccgtgggtg tegttaaagg gcgacttaga gaaccaccaa gaacgtagca 360
gaatcctcag agtgtcatac cagcatacag ccatcgctaa ctgctattta ctggtaatag 420
ggcacattgt aatctcactt aaccatactg tcggggccacc atctagccta tttctgcccac 480
gaatcaatcg tgagtgatgg acatagagaa actattagtt gagaagaaaa caagagcaact 540
aaaggtttga tattgacaaa aatctacttc gccgtcactc cataggttta ttgtctotca 600
ttagtcocaga acagcagtta cagacgtaag cttttaogca caaactacag ggttgctctt 660
tattgtatcg aaaatatggg acctgaataa gggogatttt gacgcgtcct gcccgcccat 720
tcccgatcct acggacagaa tggcaagcag tcgaogtogo cccaaacacg tcatttcgga 780
tcctcacgat ccaactaacg tgcttttaggt acctcaagca cgggtcatcg ttctcgtcgg 840
accogtcgct tgogacgaag ggctcgacga gcaaatatc cctcagacac agcccactga 900
gtttctcgcc ggatcttctc agcgggtogc gtttccgato cggtggtaga ttctgcgaag 960
cacggctctt gctaggatct gtgttagcaa cgtogtcagg tttgagcccc gtgagctcac 1200
ttactagtta aggttacgct gaaatagcct ctcaaggctc tcagctaggt aggaacacaa 1080
aaaaaaagtc ctgcccttaa caccgttgog atggcttgto ttctgcagcg tactgtcgtg 1140
gcagggcggg accgcacat ctttttcgac gccacctgt gatctgaagg cgaagatact 1200
cgaccttaat gattgaggca agagcgtaat acctcgcgct ccttagacga gtagatctcg 1260
tggaagattc ggcacacggc acacaaaaat agcttttgag atagccttca atgtaattat 1320
gtttttatat atatttacta gctgacccgg caaacgttgt gttgccttaa ataagatttc 1380
tagggaaatt ctagtgtaga aaaataacct cattcaacca cataatacct cattataacc 1440
aaaaaaaaat aatatccaaa aaataaaaat ataaaaataa tgtttggggt ggacaaccct 1500
tatcacatag gggatatgaa attagatagt agccgattct cagacctact gaacatacta 1560
ttgatacaca aataaaacca aaaaaacatg gctgaaaaat gtatagtagg tattgtatta 1620
ttaagtgtat aatctatgat gtatatgagt aagtaagaca ggagaccggc ttcgtcctca 1680
tccgtcataa aaacogcat aaaaatcaaa ccogcaaaat tataatttgc gtaattactg 1740
gtggctggtg gtaggacctt cttgtgagtc cgcgogggta ggtaccacca tctgactatt 1800
ctgccgtgaa gcagtaatgg gtttcggttt gaaggcgggg acagccgttg taactatact 1860
tgagacctta gaacttatat ctcaatgtgg gtggcgcat tttttacggt aggcagcggc 1920
ttggtctctc cctggcatt gctgaagtc ataggcgacg gttaccactc accatcaggt 1980
gggcggtatg gccgtctgcc tacaaaaatc ataaaaaaaa aataaaaaat ttacgttgta 2040
gatgtotatg ggctccagta accacttaac accaggcggg ctgtgagctc gtccacccat 2100

```

ctaagcaata aaaataaata aatagatagt tgatcagtag tggacggcg agggggggag 2160
 atcaaattga atttaaaata aacataatt aaaggaattt gaaactataa actotgaata 2220
 ataatttato gtactacaat tataatattt gattgccatc ttgcaacott attgoggatc 2280
 tgtgaataga aaaaaaaaaa aaatcgggat ggaaaaatag gggttgatcg tataagagtg 2340
 aaaattgaga gtaatatgga atttttttat ttttaagtcac gacaaaataa aaataagatc 2400
 ttgocaaaaa aattttaagtt tattattaaa ttaagtttaa caaataaaaa attgggggtg 2460
 atcgcagagg ggtgaaaatt tagggtttta tgtatttttg tatgctgtat cataaaaaaa 2520
 taaaaacaaa aaataaaaat aggggggatga aaaataaatg ttgttcgatt ctcaaccctg 2580
 gccgatatgc acgctaagat tcacaaaaat cggctcgagcc gtttcggagg agttcaatca 2640
 cgcacccctg cacacgagaa ttttatttat tagatttaga agagctgaaa gataaatcga 2700
 tattttaattt tgtaagttgt cttgatgata cattttttcg tttgtcatc tttcctgcag 2760
 ttagaacata atataaatg caaatgaaaa atagaatat aataaataat aataaataaa 2820
 taataaatat ttactaacia tcacgtacg ttaactggtc ccgtgataag ttctgaaaga 2880
 acttgtgtta caggtaaccag ataacggata taaatgtaag atttttatta tacacataca 2940
 tatatttcat atacattcat aaccctggaa aatacattta tatttatcat acaaatatct 3000
 tcccttggcg ggattcgaac ccgcgacccc cttgtgtagt gacaatgtca cttaccacta 3060
 caccctctgg cattgctggg cgacggtaac caccaccat taggtgggac atatgctcgt 3120
 ctgcctacaa gggaaataaa aaaaatatcc taatataaat tgcatattt tttttaaac 3180
 gactttcaat cacaatgaag acagattctc gtcgaagttt gtttttgaaa ctatatcaat 3240
 aacttttcat tatccgttct tcgtcttttg tcttttttc gcaaacaaaa cgaacaaaa 3300
 gttctaattc gaaagatgtt ttgtacggaa agtttgaaat agtgcttaat tgcaagtaac 3360
 gtaacaatgt tttagggttc ggtcctcaat aaattcgacc aataaaccat acaaattctt 3420
 taacattttt ttaatttat actagctgac ccggcagact tcgtgggtgc tcaatcgata 3480
 aataaaatac ctatgcttct gtataaaata aacataaaac aaacaaaag aatccgtccg 3540
 acgggagaca catcaaagga aaaacatctt ttttattttt ttacotttta aacctctct 3600
 ggacttcac aaataattta agaccaaaat tagccaaato ggtctagcat ttcgagttt 3660
 tagcgagact aacgaacagc aattcatttt tatatacaca gatttatgtt accggggtct 3720
 agtgacctaa acgacttcag ctctaacact aggctaacto aggttagta gcctggctct 3780
 agtgtttagat ttgaagtcgt ctaatgcaaa gattattgga tctgatggat ccgtaaggac 3840
 gtgtctagag cgtcgacggt gactagctcc tgctgatca ggaaaaatgt ggaaagctta 3900
 acgattttgt cacattttac ttatcacaac ttgtttttat aataattcgc ttaaatgago 3960
 agctattact taatctcgt gtggtttttg acaaaatcag cttctttaga actaaaatat 4020
 catttttttc gtaatttttt taatgaaaaa tgctctagtg ttataccttt ccaaaatcac 4080

cattaattag gtagtgttta agcttggtgt aaaaaactgc cacacgcatt tttttctcca 4140
 ctgtaggttg tagttacgog aaaacaaaat ogttctgtga aaattcaaac aaaaatattt 4200
 tttcgtaaaa acacttatca atgagtaaag taacaattca tgaataattt catgtaaaaa 4260
 aaaaatacta gaaaaggaat ttttcattac gagatgctta aaaatctgtt tcaaggtaga 4320
 gatttttoga tatttcggaa aattttgtaa aactgtaaat ccgtaaaatt ttgctaaaca 4380
 tatattgtgt tgttttggtta agtattgacc caagctatca cctoctgcag tatgtogtgc 4440
 taattactgy acacattgta taacagttcc actgtattga caataataaa acctcttcat 4500
 tgacttgaga atgtctggac agatttggtt ttgtattttt gatttacaaa tgtttttttg 4560
 gtgatttacc catccaaggo attctccagg atggttggtg catcacgcog attggcaaac 4620
 aaaaactaaa atgaaactaa aaagaaacag tttccgctgt ccggttcctc tagtgggaga 4680
 aagcatgaag taagttcttt aaatattaca aaaaaattga acgatattat aaaattcttt 4740
 aaaaatttaa aagtaagaac aataagatca attaaatcat aattaatcac attgttcatg 4800
 atcacaaatt aatttacttc atacgttgta ttgttatggt aaataaaaag attaatctct 4860
 atgtaattgt atctgtacaa tacaatgtgt agatgtttat tctatcgaaa gtaaatatcg 4920
 caaaactcga aaattttcag tataaaaagg ttcaactttt tcaaatcagc atcagttcgg 4980
 ttccaactct caagatgaga gtcaaaacct ttgtgatctt gtgctgcgct ctgcagggtga 5040
 gtttaattatt ttactattat ttcagaaggt ggccagaoga tatcacgggc cacctgataa 5100
 taagtggctg ccaaaacgca cagatatcgt aaattgtgcc atttgatttg tcacgcccgg 5160
 gggggctacg gaataaacta catttattta tttaaaaaat gaaccttaga ttatgttaact 5220
 tgtgatttat ttgogtcaaa agtaggcaag atgaatctat gtaaataacct gggcagactt 5280
 gcaatatcct atttcacogg taaatcagca ttgcaatatg caatgcata tcaacaatat 5340
 gtaaaacaat togtaaagca tcattagaaa atagacgaaa gaaattgcat aaaattataa 5400
 ccgcattatt aatttattat gatattctatt aacaattgct attgcctttt tttcgcaaat 5460
 tataatcatt ttcataacct cgaggtagca ttctgttaca ttttaataca ttggtatgtg 5520
 attataacac gagctgcccc ctgagtttct cgccagatct tctcagtggg tcgcgttacc 5580
 gatcacgtga tagattctat gaagcaactgc tcttggttag gotagtgtta gcaaattctt 5640
 tcaggttgag totgagagct cacctaccca tcggagcgta gctggaatag gctaccagct 5700
 aataggtagg gaaaacaaag ctogaaacaa gotcaagtaa taacaacata atgtgaccat 5760
 aaaaatctgt ggtgtatgag atacaattat gtactttccc aaaaatgttt acataattag 5820
 aatgttggtc aacttgccca acgcccagc tagaacattc aattattact attaccacta 5880
 ctaaggcagt atgtcctaac togttccaga tcagcgctaa cttogattga atgtgcgaaa 5940
 tttatagctc aatatttttag cacttatcgt attgatttaa gaaaaattg ttaacatttt 6000
 gtttcagtat gtcgcttata caaatgcaaa catcaatgat tttgatgagg actatttttg 6060

gagtgatgtc 6070
 <210>24
 <211>99
 <212>DNA
 <213>Bombyx mori
 <400>24
 cgcagttacg actattctcg tcgtaacgtc cgcaaaaact gtggaattcc tagaagacaa 60
 ctagttgtta aattcagagc actgcottgt gtgaattgc 99
 <210>25
 <211>28
 <212>DNA
 <213>Artificial sequence
 <400>25
 ggcgcgcgg gagaaagcat gaagtaag 28
 <210>26
 <211>26
 <212>DNA
 <213>Artificial sequence
 <400>26
 gtgcacottg agagttggaa ccgaac 26
 <210>27
 <211>24
 <212>DNA
 <213>Artificial sequence
 <400>27
 gtgcacatgg cgtgccctc ttcc 24
 <210>28
 <211>26
 <212>DNA
 <213>Artificial sequence
 <400>28
 tgtggatcct ttctogctcc ttaatc 26
 <210>29

<211>32
 <212>DNA
 <213>Artificial sequence
 <400>29
 ggatcctaata ttttaataata aaataaccct tg 32
 <210>30
 <211>35
 <212>DNA
 <213>Artificial sequence
 <400>30
 cttggcgcgc caagacgtag acgtatagcc atcgg 35
 <210>31
 <211>31
 <212>DNA
 <213>Artificial sequence
 <400>31
 cttgtcgacg acatcaactcc caaaatagtc c 31
 <210>32
 <211>33
 <212>DNA
 <213>Artificial sequence
 <400>32
 gtccgatccc gcagttacga ctattctcgt cgt 33
 <210>33
 <211>31
 <212>DNA
 <213>Artificial sequence
 <400>33
 cccaatttgg cgcgcctcaa gacatccttg a 31
 <210>34
 <211>27
 <212>DNA
 <213>Artificial sequence

<400>34	
gaatgctacc tcgagggttat gaaaatg	27
<210>35	
<211>24	
<212>DNA	
<213>Artificial sequence	
<400>35	
aacctcgagg tagcattctg ttac	24
<210>36	
<211>26	
<212>DNA	
<213>Artificial sequence	
<400>36	
ggtaccggcg cgccacgacg tagacg	26
<210>37	
<211>30	
<212>DNA	
<213>Artificial sequence	
<400>37	
gtcgacatgt cgtttacttt gaccaacaag	30
<210>38	
<211>30	
<212>DNA	
<213>Artificial sequence	
<400>38	
ggatcctttt tgacaccaga ccaactggta	33

- 54 -

CLAIMS

1. A process for producing recombinant cytokine comprising producing a gene recombinant silkworm that incorporates cytokine gene in its chromosomes, producing
5 recombinant cytokine protein in the silk glands or cocoon and silk thread of the resulting gene recombinant silkworm, and recovering the cytokine from the silk glands or cocoon and silk thread.
2. A process for producing recombinant cytokine
10 according to claim 1 wherein a cytokine gene coupled downstream from a promoter specifically expressed in silk glands is incorporated in a chromosome.
3. A process for producing recombinant cytokine according to claim 2 wherein the promoter specifically
15 expressed in silk glands is a sericin gene promoter.
4. A process for producing recombinant cytokine according to claim 2 wherein the promoter specifically expressed in silk glands is a fibroin H chain gene promoter.
20
5. A process for producing recombinant cytokine according to any of claims 1 through 4 wherein cytokine gene is incorporated in silkworm chromosomes using DNA originating in a transposon.
6. A process for producing recombinant cytokine
25 according to claim 5 wherein the cytokine gene is located between a pair of inverted terminal sequences originating in a transposon.
7. A process for producing recombinant cytokine according to claim 5 or 6 wherein the DNA originating in
30 a transposon originates in an insect.
8. A process for producing recombinant cytokine according to claim 7 wherein the transposon originates in piggyBac transposon originating in a lepidopteron.
9. A process for producing recombinant cytokine
35 according to any of claims 1 through 8 wherein the cytokine gene is interferon gene or colony stimulating factor gene.

10. A process for producing recombinant cytokine according to claim 9 wherein the interferon gene or colony stimulating factor gene is feline interferon- ω gene, human interferon- β gene or feline granulocyte colony stimulating factor gene.
11. A process for producing recombinant cytokine according to any of claims 1 through 3 wherein cytokine is extracted from cocoon and silk thread by using an aqueous solvent.
12. A gene recombinant silkworm in which a cytokine gene has been inserted into a chromosome and cytokine is produced in silk glands or cocoon and silk thread.
13. A gene recombinant silkworm according to claim 12 wherein the cytokine gene inserted into a chromosome is an interferon gene or colony stimulating factor gene.
14. A gene recombinant silkworm according to claim 13 wherein the interferon gene or colony stimulating factor gene inserted into a chromosome is feline interferon- ω gene, human interferon- β gene or feline granulocyte colony stimulating factor gene.
15. A vector for inserting an exogenous gene into silkworm chromosomes in which a cytokine gene is coupled downstream from a promoter that is specifically expressed in silk glands.
16. A vector for inserting an exogenous gene into silkworm chromosomes according to claim 15 wherein the promoter is sericin gene promoter.
17. A vector for inserting an exogenous gene into silkworm chromosomes according to claim 15 wherein the promoter is a fibroin H chain gene promoter.
18. A vector for inserting an exogenous gene into silkworm chromosomes according to any of claims 15 through 17 wherein the cytokine gene is located between a pair of inverted terminal sequences originating in a transposon.
19. A vector for inserting an exogenous gene into

- 56 -

silkworm chromosomes according to any of claims 15 through 18 wherein the cytokine gene is an interferon gene or a colony stimulating factor gene.

5 20. A vector for inserting an exogenous gene into silkworm chromosomes according to claim 19 wherein the interferon gene or colony stimulating factor gene is feline interferon- ω gene, human interferon- β gene or feline granulocyte colony stimulating factor gene.

10 21. A gene cassette for expressing an exogenous protein comprising (1) a promoter expressed in silk glands, and (2) a gene coupled downstream from (1) in which the 5' terminal portion of fibroin H chain gene is fused to the 5' side of an exogenous protein structural gene.

15 22. A gene cassette for expressing an exogenous protein comprising (1) a promoter expressed in silk glands, and (2) a gene coupled downstream from (1) in which the 3' terminal portion of fibroin H chain gene is fused to the 3' side of an exogenous protein structural
20 gene not containing a stop codon, or a gene cassette for expressing an exogenous protein comprising (1) a promoter expressed in silk glands, and (2) a gene coupled downstream from (1) in which an exogenous protein structural gene is fused to the 3' side of the 3'
25 terminal portion of fibroin H chain gene.

23. A gene cassette for expressing an exogenous protein comprising (1) a promoter expressed in silk glands, and (2) a gene coupled downstream from (1) in which the 5' terminal portion of fibroin H chain gene is
30 fused to the 5' side of an exogenous protein structural gene not containing a stop codon, and in which the 3' terminal portion of fibrin H chain gene is fused to the 3' side of the structural gene.

24. A gene cassette according to claim 21 or 23
35 wherein the 5' terminal portion of the fibroin H chain gene contains a first exon, first intron and a portion of a second exon of fibroin H chain gene.

- 57 -

25. A gene cassette according to claim 24 wherein the portion where the first exon and second exon of the fibroin H chain gene are joined is a secretion signal gene region of fibroin H chain gene.

5 26. A gene cassette according to claim 25 wherein the promoter expressed in silk glands of (1) and the 5' terminal portion of fibroin H chain gene coupled downstream from (1) are the DNA shown in SEQ. ID No. 22 and SEQ. ID No. 23.

10 27. A gene cassette according to claim 22 or 23 wherein the 3' terminal portion of the fibroin H chain gene contains at least one codon that encodes cysteine.

15 28. A gene cassette according to claim 27 wherein the 3' terminal portion of the fibroin H chain gene is the DNA shown in SEQ. ID No. 24.

20 29. A gene cassette according to any of claims 21 through 28 wherein the promoter expressed in silk glands is at least one promoter selected from fibroin H chain gene promoter, fibroin L chain gene promoter and sericin gene promoter.

25 30. A gene cassette according to any of claims 21 through 29 wherein at least one poly A addition region selected from a poly A addition region of fibroin H chain gene, a poly A addition region of fibroin L chain gene and a poly A addition region of sericin gene is present downstream from a gene cassette for expressing an exogenous protein according to any of claims 21 through 29.

30 31. A gene cassette for inserting a gene into chromosomes of insect cells comprising inverted repetitive sequences of a pair of piggyBac transposons present on both sides of a gene cassette for expressing an exogenous protein according to any of claims 21 through 30.

35 32. An expression vector for insect cells that contains a gene cassette for expressing an exogenous protein according to any of claims 21 through 31.

- 58 -

33. A gene insertion vector for insect cells that contains a gene cassette for inserting a gene into chromosomes of insect cells according to claim 31.

5 34. A process for producing an exogenous protein comprising inserting a vector for insect cells according to claim 32 or 33 into insect cells.

35. A process for producing an exogenous protein according to claim 34 wherein the insect cells originate in a lepidopteron.

10 36. A process for producing an exogenous protein according to claim 35 wherein the insect cells originate in silkworm moths (*Bombyx mori*).

15 37. A process for producing an exogenous protein according to claim 36 wherein the insect cells are silk gland cells of silkworm moths (*Bombyx mori*).

20 38. A process for producing an exogenous protein comprising producing a recombinant silkworm in which a gene cassette for expressing an exogenous protein according to any of claims 21 through 31 is inserted into a chromosome using a gene insertion vector for insect cells according to claim 33 and the DNA transfer activity of piggyBac transposase, producing exogenous protein in the silk glands or cocoon and silk thread of the resulting recombinant silkworm, recovering the exogenous protein from the silk glands or silk and cocoon thread.


25 39. A process for producing an exogenous protein according to claim 38 wherein the recombinant silkworm, in which the gene cassette for expressing an exogenous protein has been inserted into a chromosome, is produced by simultaneously micro-injecting the gene insertion vector for insect cells and DNA or RNA that produces the piggyBac transposase into silkworm eggs.

30 40. A recombinant silkworm in which a gene cassette for expressing an exogenous protein according to any of claims 21 through 31 has been inserted into a chromosome, and which has the ability to produce the exogenous protein in silk glands or silk thread.

- 59 -


41. Silk thread containing an exogenous protein
produced by a recombinant silkworm according to claim 40.

Fig.1



SerP: SERIGIN-1 GENE PROMOTER
 IFN: FELINE INTERFERON- ω GENE
 BGHpA: BOVINE GROWTH HORMONE POLY A
 A3P: A3 PROMOTER
 GFP: GREEN FLUORESCENCE PROTEIN
 SV40pA: SV40 POLY A
 ITR: INVERTED TERMINAL SEQUENCE

Fig.2



FibHP: FIBROIN H CHAIN GENE PROMOTER
 IFN: FELINE INTERFERON- ω GENE
 BGHPA: BOVINE GROWTH HORMONE POLY A
 A3P: A3 PROMOTER
 GFP: GREEN FLUORESCENCE PROTEIN
 SV40pA: SV40 POLY A
 ITR: INVERTED TERMINAL SEQUENCE

Fig.3




Fig.4




Fig.5




Fig.6




Fig.7




Fig.8




Fig.9




Fig.10





Fig.11

Fibroin H promoter 0.3kbp.






Fig.13




Fig.14

FROM FIG. 13




Fig.15




Fig.16




Fig.17




Fig.18




Fig.19

